

(19)



Eur päisches Patentamt
Eur pean Patent Office
Offic européen des brevets



(11)

EP 0 967 278 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
29.12.1999 Bulletin 1999/52

(51) Int Cl.⁶: **C12N 15/29, C12N 15/82,
C07K 14/415, A01H 5/00,
C12N 15/11, C07K 16/16**

(21) Application number: **99305077.2**

(22) Date of filing: **28.06.1999**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **26.06.1998 JP 18006598
24.06.1999 JP 17904399**

(71) Applicant: **Mitsui Chemicals, Inc.
Tokyo 100-6070 (JP)**

(72) Inventors:
• **Yoshida, Nobumasa
Mobara-shi, Chiba 297-0017 (JP)**

- **Kato, Yoshihiro
Mobara-shi, Chiba 297-0029 (JP)**
- **Takahashi, Shigeru
Mobara-shi, Chiba 297-0022 (JP)**
- **Yanai, Yukihiro
Mobara-shi, Chiba 297-0026 (JP)**
- **Hiratsuka, Junzo
Mobara-shi, Chiba 297-0017 (JP)**
- **Miwa, Tatsushi
Nerima-ku, Tokyo 176-0024 (JP)**

(74) Representative:
**Holdcroft, James Gerald, Dr. et al
Graham Watt & Co.,
Riverhead
Sevenoaks, Kent TN13 2BN (GB)**

(54) **Flowering regulating gene and its use**

(57) Flowering regulating genes of plants and methods for controlling plant flowering are provided. The flowering time can be modified in comparison with wild

type plants by enhancing or inhibiting the expression of the flowering regulating gene. Transgenic plants in which the expression of the flowering regulating gene is regulated is also provided.

EP 0 967 278 A2

Description**FIELD OF THE INVENTION**

5 [0001] The present invention relates to genes for floral regulation of plants and to methods for controlling plant flowering by regulating the expression of said gene. The present invention also relates to transgenic plants whose flowering time is modified in comparison with wild type plants by regulating the expression of said gene and to methods for generating said transgenic plants.

10 **BACKGROUND OF THE INVENTION**

[0002] In order to resolve the worldwide food problem, developing technology for increasing the yield of food using biotechnology has been desired. Grain, which is one of main crops, is seed of plants and some vegetables are fruits of plants. For productivity increase of these plants, floral regulation for controlling growth of plants is an important key technology. On the other hand, flowering inhibition of vegetables, whose vegetative organs such as leaves or roots are marketed, prevents vegetative organs from stopping their growth and often increases their productivity. In addition, for many crops the suitable cultivating places are limited because of their species specificity of hereditary flowering behavior depending on environment. Modification of these properties by flowering regulation can expand the suitable cultivating places.

20 [0003] In molecular genetic studies using model plants such as *Arabidopsis thaliana* and *Antirrhinum majas*, many genes involved in identity determination of floral meristems or morphogenesis of floral organs have been isolated. Among these genes LEAFY and APETALA-1 genes are known to be forcedly expressed in the host plant *Arabidopsis* or poplar when introduced into these plants, thereby flowering the plants earlier. Since these genes are not fundamentally involved in floral budding (the transition from vegetative growth to reproductive growth), the use of these genes alone cannot arbitrarily regulate flowering. If the function of these genes is inhibited, the shape of inflorescence is changed, which is obvious from the phenotype of the mutants, and flowering cannot be regulated.

25 [0004] The embryonic flower mutant of *Arabidopsis*, in which flowering occurs immediately after germination, is known (Sung et al. (1992), Science, vol.258: p1645-). In this mutant, the function of a gene that maintains vegetative growth for a certain period of time after germination is thought to be lost. The flowering of wild-type *Arabidopsis* is thought to be inhibited by the expression of this gene. Although the approximate location of this gene on the chromosome is reported (Yang et al. (1995), Dev. Biol., vol.169: p421-), the result is far from helping the isolation of the gene and the gene has not yet been isolated.

35 **SUMMARY OF THE INVENTION**

[0005] An objective of the present invention is to isolate a gene for floral regulation (flowering regulating gene) and to provide a transgenic plant into which the gene is introduced. If a fundamental gene that regulates flowering is isolated, flowering time can be freely controlled by artificially regulating this gene.

40 [0006] The present inventors have succeeded in isolating mutant *Arabidopsis* that exhibits flowering immediately after germination because the function of the flowering regulating gene is lost and in identifying a single gene, which was mutated, in a wide region of the chromosome and isolating it. Furthermore, the present inventors have confirmed that this gene has flowering inhibiting function by introducing the gene into *Arabidopsis* and expressing it. Based on these findings, the present inventors have completed the present invention.

45 [0007] Moreover, the present inventors have discovered that the flowering regulating gene isolated from any kind of plant by hybridization or PCR technique based on the sequence of *Arabidopsis* flowering regulating gene has the function that complements the mutation of the *Arabidopsis* super early flowering mutant, inhibits flowering, and induces normal differentiation of stems and leaves.

50 [0008] Thus, the present invention relates to novel flowering regulating genes that exist extensively in plants, proteins with flowering regulating activity encoded by said genes, transgenic plants in which the expression of said gene is modified, methods for generating these plants, and methods for controlling the flowering time of plants by regulating the expression of said genes. More specifically, the present invention relates to

(1) a DNA encoding a protein having flowering regulating activity, wherein said DNA selected from the group consisting of:

55

- i) a DNA encoding a protein comprising the amino acid sequence of SEQ ID NO: 1;
- ii) a DNA encoding a protein comprising the amino acid sequence substantially identical to that of SEQ ID NO: 1;
- iii) a DNA hybridizing a DNA encoding the protein comprising the amino acid sequence of SEQ ID NO: 1;

- iv) a DNA encoding a protein comprising amino acid sequences showing 50% or more and 60% or more homology with amino acids 278 to 348 and 465 to 607, respectively, of the amino acid sequence of SEQ ID NO: 1;
- v) a DNA encoding a protein comprising the amino acid sequence of SEQ ID NO: 8.
- 5 vi) a DNA encoding a protein comprising the amino acid sequence substantially identical to that of SEQ ID NO: 8.
- vii) a DNA hybridizing a DNA encoding the protein comprising the amino acid sequence of SEQ ID NO: 8; and
- viii) a DNA encoding a protein comprising amino acid sequences showing 50 % or more and 60 % or more homology with amino acids 282 to 352 and 450 to 592, respectively, of the amino acid sequence of SEQ ID
- 10 NO: 8. a DNA encoding a protein having flowering regulating activity, wherein said protein comprises the amino acid sequence of SEQ ID NO: 1;
- (2) the DNA of (1), wherein said DNA of i) comprises the coding region of the nucleotide sequence of SEQ ID NO: 2;
- (3) the DNA of (1), wherein said DNA of v) comprises the coding region of the nucleotide sequence of SEQ ID NO: 9;
- 15 (4) the DNA of any one of (1) to (3), encoding a protein having a zinc finger structure;
- (5) a protein having flowering regulating activity, encoded by the DNA of any one of (1) to (4);
- (6) the protein of (5), comprising the amino acid sequence of SEQ ID NO: 1 or 8;
- (7) a recombinant double-stranded DNA molecule comprising an expression cassette comprising the DNA of any one of (1) to (4);
- 20 (8) a recombinant double-stranded DNA molecule comprising an expression cassette comprising the following constituent elements of i) to iii),
- i) a promoter that can transcribe in plant cells,
- ii) the DNA of any one of (1) to (4) or a part of it fused to said promoter in sense or antisense direction, and
- 25 selectively, and
- iii) a signal involved in transcription termination of RNA molecules and polyadenylation, wherein the signal functions in plants;
- (9) a transformant into which the recombinant double-stranded DNA molecule of (7) is introduced ;
- 30 (10) a transgenic plant cell into which the recombinant double-stranded DNA molecule of (8) is introduced;
- (11) a method for producing a protein of (5) or (6), wherein the method comprises
- (a) cultivating a transformant of (9) and
- (b) recovering a recombinant protein from said transformant or the culture supernatant of it;
- 35 (12) a transgenic plant comprising transgenic plant cells of (10);
- (13) a method for producing a transgenic plant of (12), wherein said method comprises
- (a) introducing the recombinant double-stranded DNA molecule of (8) into plant cells and
- 40 (b) regenerating said plant cells;
- (14) a DNA encoding an antisense RNA complementary to a transcription product of a DNA of any one of (1) to (4);
- (15) a method for regulating the flowering time of a plant, wherein said method comprises introducing the whole or a part of a DNA of any one of (1) to (4) or the whole or a part of a DNA of (14) into a plant and expressing it,
- 45 thereby changing the activity of a flowering regulating protein; and
- (16) an antibody that binds to a protein of (5) or (6).

BRIEF DESCRIPTION OF THE DRAWINGS

- 50 [0009] Figure 1 shows the location of the DNA clones of the chromosome region containing *Arabidopsis* "MPC1" gene and the markers 16EB53 and z11-1. In the figure the unfilled arrow shows the position and direction of "MPC1" gene and 11K22, 22K2, 19A20, and 20I12 show DNA clones.

DETAILED DESCRIPTION OF THE INVENTION

- 55 [0010] "An expression cassette" used herein means a DNA molecule comprising a gene and constituent elements essential for the expression of the gene. Typically, it is a DNA molecule comprising (i) a promoter to express a structural gene in a host, (ii) the structural gene, and, if necessary, (iii) a terminator. The promoter varies depending on the host.

For example, in order to produce a recombinant protein in a microorganism, a promoter functioning in the microorganism is used. For generating a transgenic plant, a promoter functioning in plant cells is used. An example of "a recombinant double-stranded DNA molecule comprising an expression cassette" is typically a vector comprising an expression cassette.

5 [0011] The present invention provides novel proteins regulating the flowering of plants and DNAs encoding said proteins. The nucleotide sequences of the cDNA and the genomic DNA of *Arabidopsis*-derived "MPC1", which has been isolated by the present inventors, are shown in SEQ ID NO: 2 and 3, respectively. The amino acid sequence of *Arabidopsis*-derived "MPC1" protein encoded by the cDNA or genomic DNA is shown in SEQ ID NO: 1. The nucleotide sequences of the cDNA and the genomic DNA of rice-derived "Os-MPC1", which has been isolated by the present
10 inventors, are shown in SEQ ID NO: 9 and 10, respectively. The amino acid sequence of rice-derived "Os-MPC1" protein encoded by the cDNA or genomic DNA is shown in SEQ ID NO: 8.

[0012] The mutation of *Arabidopsis*-derived "MPC1" gene eliminates normal flowering regulating ability of plants and leads plant to flowering immediately after germination (super early flowering mutation). The present inventors have found that guanine is substituted with adenine at the nucleotide 5039 of "MPC1" genomic DNA (SEQ ID NO: 3) in
15 "mpc1" mutant plants (Example 1). Since the C-terminal amino acid residues from 541 of "MPC1" protein are not translated by this base substitution, the deletion of the amino acid sequence after this mutation point is thought to diminish the normal flowering regulating function of "MPC1" protein. In other words, this deletion inhibits flowering function and leads plants to super early flowering. This phenomenon has been induced by introducing and expressing an antisense DNA in plants, thereby inhibiting "MPC1" protein expression (Example 2). Furthermore, the cDNA of rice-
20 derived "Os-MPC1", which shows significant homology with said *Arabidopsis* cDNA, also complement the super early flowering mutation of *Arabidopsis* (Example 5). It is therefore thought that these proteins exist widely in plants and regulate the flowering time.

[0013] Many plants including *Arabidopsis* vegetatively grow for a certain period of time after germination, flower, and reproductively grow. The genes of the present invention are essential to maintain vegetative growth, and to regulate
25 the transition from vegetative growth to reproductive growth. In other words, the expression level of this gene regulates flowering. Therefore, the flowering time of plants can be changed by artificially regulating the expression of the genes of the present invention, which leads to productivity increase of useful plants.

[0014] DNAs used in this invention are not limited to DNAs encoding *Arabidopsis*-derived "MPC1" or rice-derived "Os-MPC1" protein mentioned above. Other DNAs encoding proteins functionally equivalent to these proteins can also
30 be used.

[0015] An example of these DNAs is a DNA encoding a protein having an amino acid sequence substantially identical to that of *Arabidopsis*-derived "MPC1" or rice-derived "Os-MPC1" protein, whose amino acid sequence is shown in SEQ ID NO: 1 or 8. "An amino acid sequence substantially identical" used herein means a sequence in which changes such as deletion, substitution, addition, and/or insertion have occurred at one or more amino acid residues of the control
35 amino acid sequence, and an amino acid sequence constitutes a protein having flowering regulating activity as the protein comprising the control amino acid sequence. Changes such as deletion, substitution, and addition can be performed at several amino acid residues, for example, by site-directed mutagenesis (Kunkel et al. (1985), Proc. Natl. Acad. Sci. USA, vol.82: p488-). Mutations of amino acids can also occur spontaneously.

[0016] Comparing amino acid sequences of proteins having flowering regulating activity of *Arabidopsis* and rice, high homology is found particularly in the region comprising zinc finger motifs and the region comprising an acidic amino acid cluster at the C-terminus. A zinc finger or zinc finger structure is a structure in which a part of a protein folds chelating zinc (Zn) to construct a protruding structure like a finger, and is thought to play an important role when the protein binds to a nucleic acid or other protein (Roosenfeld et al. (1993), J. Biomol. Struct. Dyn., Vol.11: p557-). An amino acid sequence that can form a zinc finger structure is called a zinc finger motif, several types of which are known.
40 Zinc fingers of Cys2-His2 (C2H2) type are found at the amino acids 306 to 327 of the *Arabidopsis*-derived "MPC1" protein and the amino acids 310 to 331 of rice-derived "Os-MPC1" protein. These motifs can be identified by, for example, a program such as "MOTIF" of "GenomeNet" (<http://www.genome.ad.jp/>), which is provided by Institute for Chemical Research, Kyoto University through the internet.

[0017] Acidic amino acid clusters are found in some kinds of transcription regulating proteins and sometimes play an important role in activating transcription (T. Tamura (1995), Mechanism of Transcriptional Regulation, Experimental Medicine Bioscience, Yodosha). These acidic amino acid clusters are found at amino acids 503 to 520 of *Arabidopsis*-
45 derived "MPC1" protein and amino acids 488 to 505 of rice-derived "Os-MPC1" protein. These regions comprising a zinc finger motif or acidic amino acid cluster are likely to play an important role in flowering regulation of plants, and it is expected that high homology is kept in these regions of flowering regulation related proteins derived from plants
50 other than *Arabidopsis* and rice.

[0018] The proteins having amino acid sequences substantially identical to that of *Arabidopsis*-derived "MPC1" or rice-derived "Os-MPC1" protein shown in SEQ ID NO: 1 or 8 are preferably those comprising amino acid sequences substantially identical to the regions of the above sequences comprising a zinc finger motif and a C-terminal acidic
55

amino acid cluster.

[0019] A specific example thereof is a protein having flowering regulating activity, wherein the protein comprises amino acid sequences showing 50% or more and 60% or more homology with amino acids 278 to 348 and 465 to 607, respectively, of the amino acid sequence of SEQ ID NO: 1 (*Arabidopsis*), or a protein having flowering regulating activity, wherein the protein comprises amino acid sequences showing 50% or more and 60% or more homology with amino acids 282 to 352 and 450 to 592, respectively, of that of SEQ ID NO: 8 (rice).

[0020] Whether a protein has flowering regulating activity or not can be evaluated by, for example, introducing a DNA encoding said protein into super early flowering mutant plants. For example, a DNA encoding a test protein are introduced into super early flowering mutant plants such as "mpc1" *Arabidopsis* mutant, and expressed. The introduced DNA is judged to encode a protein having flowering regulating activity if it complements super early flowering mutant and differentiates normal stems and leaves as shown in Example 5. These DNA are thought to encode proteins having the same function as *Arabidopsis*-derived "MPC1" or rice-derived "Os-MPC1" protein (SEQ ID NO: 1 or 8, respectively).

[0021] In addition, other DNAs encoding proteins functionally equivalent to *Arabidopsis*-derived "MPC1" or rice-derived "Os-MPC1" protein can be screened by hybridization technique using the whole or a part of the DNA sequence encoding the amino acid sequence of SEQ ID NO: 1 or 8 as a probe (Southern (1975), J. Mol. Biol., vol.98: p503; Sambrook et al. (1989), Molecular Cloning, Cold Spring Harbor Laboratory Press). Partial sequences of "MPC1" or "Os-MPC1" used as probes are at least fourteen or more nucleotide sequences. For example, GenelImage system (Amersham) can be used for hybridization. In accordance with the protocol attached to the product, test DNAs are incubated overnight with labeled probes, and those that hybridizes with the probes can be screened by washing at 50 °C with 6xSSC and 0.1% SDS. Alternatively, DNAs encoding proteins functionally equivalent to *Arabidopsis*-derived "MPC1" or rice-derived "Os-MPC1" protein can be isolated from other plants by PCR technique using oligonucleotides specifically hybridizing with the DNA encoding the amino acid sequence constituting "MPC1" or "Os-MPC1" protein as primers (K. Shimamoto & T. Sasaki (1995), Protocols of PCR Experiments for Plants, Cell Engineering SUPPLEMENT, Plant Cell Engineering Series 2, Shujunsha).

[0022] Flowering regulating proteins encoded by DNAs obtained by such hybridization or PCR technique are thought to have high homology with *Arabidopsis*-derived "MPC1" or rice-derived "Os-MPC1" protein. The term "high homology" means 45% or more, preferably 60% or more, more preferably 75% or more, still more preferably 90% or more, and most preferably 95% or more homology with at least one amino acid sequence of these proteins. The homology may possibly become 45% or less when plural amino acid residues of the amino acid sequence encoded by the isolated DNA are deleted, added, or replaced. Even in this case, the DNA can encode a protein having the region essential for the function of flowering regulating proteins and having the equivalent flowering regulating activity. As mentioned above, it is important for the protein to exhibit the flowering regulating function- that high homology exists, in particular, in regions comprising a zinc finger motif region and a C-terminal acidic amino acid cluster region.

[0023] The homology between two or more genes in terms of the nucleotide sequences or the amino acid sequences of the proteins encoded by the genes can be determined using software for gene analysis, for example, DNASIS (Hitachi Software Engineering). In the software, the programs "Homology Plot," which plots homology as two-dimensional image, and "Maximum Matching," in which sequences are aligned considering gaps, are available for calculating homology between two genes (Needleman, S. B. et al. (1970), J. Mol. Biol., vol.48: p443-). The "Multialignment" program aligns three or more kinds of sequences to clarify the homologous regions (Waterman, M. S. (1986), Nucleic Acids Research, vol.14: 9095-).

[0024] Examples of plants from which the DNAs of the present invention are isolated by hybridization or PCR technique include corn, wheat, barley, rye, potato, tobacco, sugar beet, sugarcane, rape seed, soybean, sunflower, cotton, orange, grape, peach, pear, apple, Japanese apricot, tomato, Chinese cabbage, cabbage, Japanese radish, carrot, pumpkin, cucumber, melon, parsley, orchid, chrysanthemum, lily, saffron, pine, eucalyptus, acacia, poplar, Japanese cedar, Japanese cypress, bamboo, and yew, in addition to *Arabidopsis* and rice, but are not limited thereto. The present inventors have succeeded in isolating a flowering regulating gene encoding a protein substantially the same as *Arabidopsis*-derived "MPC1" or rice-derived "Os-MPC1" protein from sugar beet using hybridization or PCR technique mentioned above (Example 6).

[0025] Flowering regulating proteins of the present invention can be produced as recombinant proteins or natural proteins. Recombinant proteins can be expressed with, for example, the expression system using *E. coli* as a host, as fusion proteins to glutathione S-transferase (Smith, D. B. et al. (1988), Gene vol.67: p32-) or as fusion proteins with histidine-tag (Nakamura et al. (1998), Protocols of Protein Experiments for Plants, Cell Engineering SUPPLEMENT, Plant Cell Engineering Series 9, Shujunsha). The desired protein expressed as a fusion protein in *E. coli* is isolated by purifying the fusion protein by affinity chromatography with glutathione or metal ions as ligands and cutting out the desired protein by an appropriate protease treatment. Natural proteins can be produced by known methods for preparing proteins from plants (Nakamura et al. (1998), Protocols of Protein Experiments for Plants, Cell Engineering SUPPLEMENT, Plant Cell Engineering Series 9, Shujunsha).

[0026] Using recombinant or natural flowering regulating proteins prepared by the method mentioned above, poly-

clonal or monoclonal antibodies against them can be generated (Nakamura et al. (1998), *Protocols of Protein Experiments for Plants*, Cell Engineering SUPPLEMENT, Plant Cell Engineering Series 9, Shujunsha). Polyclonal antibodies can be generated by, for example, the method below. A laboratory animal such as a mouse is immunized with the prepared protein or its partial fragments mixed with appropriate adjuvant by intraperitoneal or subcutaneous injection. Additional immunization is then performed 2 to 10 times every one to four week, preferably every one or two week. After the fourth week, the blood is collected, serum is obtained to serve as antibody, and the antibody titer is measured by, for example, western blotting. The obtained antibody can be used in various experiments.

[0027] Monoclonal antibodies can be produced by fusing myeloma cells and the spleen cells obtained from the laboratory animal such as a mouse immunized by the method mentioned above and cloning the hybridoma producing the desired antibody. The hybridoma is cultivated in an appropriate medium to obtain the desired monoclonal antibody from the culture supernatant. A large amount of antibody can be obtained when hybridoma is cultivated in ascites. For example, hybridoma is transplanted into a nude mouse and allowed to grow. The monoclonal antibody produced in ascites of said animal is then collected.

[0028] Plant flowering regulation of the present invention can be performed by enhancing or inhibiting the expression of DNAs encoding the flowering regulating proteins mentioned above in target plants. Specifically, transgenic plants are generated by introducing said DNA or the antisense DNA against said DNA to the target plant. The DNA or the antisense DNA can be placed under the control of an appropriate inducible promoter to subtly regulate the degree of activation or inhibition of flowering and flowering time.

[0029] These DNAs can be expressed by introducing, into plant cells, a recombinant double-stranded DNA molecule comprising an expression cassette comprising (i) a promoter that is transcribed in plant cells, (ii) the whole or a part of the DNA encoding a flowering regulating protein of the present invention fused at the downstream of the promoter in sense or antisense direction, and if necessary, (iii) a terminator sequence fused at the downstream of the DNA, which comprises a polyadenylation site essential for stabilizing the transcript. "A part of the DNA encoding a flowering regulating protein" used herein means a part of the DNA encoding a complete flowering regulating protein that regulates flowering when it is expressed in plant cells. The present invention includes these recombinant double-stranded DNA molecules. The recombinant double-stranded DNA molecules can have DNA sequences essential to transfer the molecule to host plant cells or to maintain it in the host cells at its 5'- and/or 3'-end as well as constituent elements described above.

[0030] An expression cassette can comprise a promoter to express constitutively or inducibly the DNA encoding the inserted flowering regulating protein of the present invention. Examples of promoters for constitutive expression are 35 S promoter of cauliflower mosaic virus (Odell et al. (1985), *Nature*, vol.313: p810-) and rice actin promoter (Zhang et al. (1991), *Plant Cell*, vol.3: p1155-). Examples of promoters for inducible expression are promoters known to express by external factors such as infection or invasion of fungi, bacteria, or virus, low or high temperature, dryness, irradiation of ultraviolet rays, contacting with specific compounds. Examples of these promoters are rice chitinase gene promoter (Xu et al. (1996), *Plant Mol. Biol.*, vol.30: p387-) and tobacco PR protein gene promoter (Ohshima et al. (1990), *Plant Cell*, vol.12: p95-), both of which are induced by infection or invasion of fungi, bacteria, or virus, rice "lip19" gene promoter that is induced by low temperature (Aguan et al. (1993), *Mol. Gen. Genet.*, vol.240: p1-), *Arabidopsis* "HSP18.2" gene promoter that is induced by high temperature (Yoshida et al. (1995), *Appl. Microbiol. Biotechnol.*, vol.44(3-4): p466-), rice "rab" gene promoter that is induced by dryness (Yamaguchi-Shinozaki et al. (1990), *Plant Mol. Biol.*, vol.14(1): p29-), parsley chalcone synthase gene promoter that is induced by ultraviolet rays (Schulze-Lefert et al. (1989), *EMBO J.*, vol.8: p651-), and corn alcohol dehydrogenase gene promoter that is induced under anaerobic conditions (Walker et al. (1987), *Proc. Natl. Acad. Sci. USA* vol.84: p6624-). Besides, rice chitinase gene promoter and tobacco PR protein gene promoter are induced by specific compound such as salicylic acid, and rice "rab" gene promoter by sprinkling of a plant hormone abscisic acid.

[0031] Various cloning vectors comprising the replication origin of *E. coli* and a marker gene for screening transformed bacterial cells are available to introduce the recombinant DNA molecules into plants. Examples of these vectors include pBR322, pUC series, and M13mp series. A desired sequence can be introduced into a vector at an appropriate restriction enzyme site. A plasmid DNA obtained can be characterized by restriction endonuclease cleavage site analysis, gel electrophoresis, and other biochemical-molecular biological methods. Once the plasmid DNA is prepared, it can be cleaved and ligated with another DNA. The sequence of the plasmid DNA can be cloned into the same plasmid or other plasmids.

[0032] When the whole of a DNA encoding a flowering regulating protein of the present invention, for example, the whole region of *Arabidopsis*-derived "MPC1" cDNA, shown in SEQ ID NO: 2, is ligated at the downstream of a promoter in sense direction, expression of the flowering regulating gene can be expressed constitutively or inducibly depending on the property of the promoter used. Then, the activity of the flowering regulating protein in plant cells constitutively or inducibly increases, and consequently, delay or inhibition of flowering can constitutively or inducibly be caused in plants.

[0033] When the whole or a part of a DNA encoding a flowering regulating protein, for example, the whole or a part

region of *Arabidopsis*-derived "MPC1" cDNA, shown in SEQ ID NO: 2, is ligated at the downstream of a promoter mentioned above in antisense direction, the antisense RNA complementary to the transcript of "MPC1" cDNA can be constitutively or inducibly expressed depending on the property of the promoter used. The expression of a flowering regulating protein of the present invention is constitutively or inducibly inhibited in plant cells, and consequently, flowering can be enhanced constitutively or inducibly in plants. Antisense DNAs used do not have to encode the antisense RNAs completely complementary to the transcript of endogenous flowering regulating protein gene as long as it can inhibit the expression of endogenous flowering regulating protein.

[0034] In plants, when a gene is ligated in sense direction at the downstream of a promoter causing constitutive and strong expression, the expression of both of the gene introduced and the corresponding endogenous gene is sometimes inhibited (Montgomery (1998), Trends Genet., 14, 255-). This phenomenon is called co-suppression. When a flowering regulating gene of the present invention, for example, the whole region of *Arabidopsis*-derived "MPC1" cDNA, shown in SEQ ID NO: 2, is ligated in sense direction at the downstream of 35 S promoter, the expression of the endogenous flowering regulating protein in plant cells can be inhibited by co-suppression to enhance flowering in plants.

[0035] Moreover, when a part of a DNA encoding a flowering regulating protein of the present invention, for example, a part region of "MPC1" cDNA, shown in SEQ ID NO: 2, is ligated at the downstream of a promoter mentioned above in sense direction, an incomplete flowering regulating protein can be constitutively or inducibly expressed depending on the property of the promoter used. The incomplete flowering regulating protein that constitutively or inducibly accumulates in cells can inhibit the normal function of the flowering regulating protein, thereby enhancing flowering constitutively or inducibly in plants.

[0036] Both dicotyledon and monocotyledon can be used as target plants for generating plants whose flowering behavior is changed in comparison with wild type plants. Particularly important plants are grain (for example, rye, wheat, corn, barley, and rice), fruits (for example, orange, grape, peach, pear, apple, and Japanese apricot), vegetables (for example, tomato, Chinese cabbage, cabbage, Japanese radish, carrot, pumpkin, potato, cucumber, melon, and parsley), ornamental plants (for example, orchid, chrysanthemum, lily, and saffron), other industrial crops of economical importance (for example, tobacco, sugar beet, rape seed, soybean, sunflower, and cotton), and trees that require a long period till flowering (for example, eucalyptus, acacia, and poplar, which are used as wood pulp, and cedar, Japanese cypress, pine, bamboo, and yew, which are used as lumber).

[0037] Various methods can be used for introducing expression cassettes into plant host cells. Examples thereof are transformation of plant cells with T-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as a transformation mediator, direct introduction into protoplast (infection method, electroporation method, etc.), and particle gun method, but are not limited thereto.

[0038] Direct introduction into protoplast needs no special vectors. For example, simple plasmids such as pUC derivatives can be used. Some methods for introducing a desired gene into plant cells need other DNA sequences. For example, when a Ti or Ri plasmid is used to transform plant cells, at least right side sequence or usually both side sequences at the T-DNA region of Ti or Ri plasmid should be connected adjacent to a gene to be introduced.

[0039] When *Agrobacterium* is used to transform plant cells, an expression cassette to be introduced should be cloned in a special plasmid, an intermediate vector or binary vector. An intermediate vector is not replicated in *Agrobacterium*. An intermediate vector is transferred into *Agrobacterium* with a helper plasmid or by electroporation. Having regions homologous to T-DNA sequence, an intermediate vector is integrated into Ti or Ri plasmid of *Agrobacterium* by homologous recombination. *Agrobacterium* used as a host has to comprise vir region. Usually, Ti or Ri plasmid comprises vir region and can transfer T-DNA into plant cells by its function.

[0040] In contrast, since a binary vector can be replicated and maintained in *Agrobacterium*, if it is introduced into *Agrobacterium* with a helper plasmid or by electroporation, T-DNA on a binary vector can be transferred into plant cells by the function of vir region of the host. The present invention also includes intermediate vectors or binary vectors thus obtained, and microorganisms such as *E. coli* or *Agrobacterium* comprising them.

[0041] Transformed plant cells can be regenerated to a plant. The method for regeneration depends on the kind of the plant cells. Examples thereof are the methods of Fujimura et al. (Fujimura et al. (1995), Plant Tissue Culture Lett., vol.2: p74-) for rice, Shillito et al. (Shillito et al. (1989), Bio/Technology, vol.7: p581-) for corn, Visser et al. (Visser et al. (1989), Theor. Appl. Genet., vol.78: p594-) for potato, and Akama et al. (Akama et al. (1992), Plant Cell Rep., vol. 12: p7-) for *Arabidopsis*. In plants generated by these methods or plants obtained from their vehicles for reproduction (for example, seeds, tubers, cuttings), the flowering regulating protein expression of the present invention changes in comparison with wild type plants, which changes the flowering behavior. The present invention includes transgenic plants thus obtained.

[0042] The present invention provides a novel gene that inhibits flowering of plants. When this gene is introduced into other plants and expressed in the plants, it can inhibit or enhance flowering of the plants.

[0043] Various cultivars of grain and vegetable that matures earlier or later than usual can be generated by regulating flowering, which produces such an agriculturally important value as expansion of the suitable cultivation place, increase of yield, and supply of crops with high value added. In particular, though the deterioration of quality by bolting and

flowering is a problem in leaf and stem vegetables such as Chinese cabbage and root vegetables such as Japanese radish, the type of cultivation is limited at present. Therefore, flowering inhibition will bring a great effect such as the expansion of the cultivation season and the suitable cultivation place. Arbitrary flowering regulation will also be considerably useful if it is applied to a cultivar having superior characteristics such as good taste or strong disease resistance. Furthermore, enhancing the floral budding of fruits will increase their productivity or change the flowering time, which enables the production and shipment of fruits out of season. In addition, inhibiting flowering of wood will not only enhance alternation of generations by shortening the period required for flowering but also enhance vegetative growth or suppress allergy induction in humans caused by scatter of pollen, which is economically and socially significant.

[0044] The present invention is illustrated in detail below with reference to Examples, but is not to be construed as being limited thereto. Besides, methods for general gene recombination such as cleavage and ligation of DNAs, transformation of *E. coli*, determination of nucleotide sequences of genes, hybridization were performed, unless otherwise mentioned, based on manuals attached to commercial reagents and apparatus or laboratory books, for example, "Molecular Cloning" (Sambrook et al. (1989), Cold Spring Harbor Laboratory Press). In addition, cultivation of *Arabidopsis* using agar medium or soil, mating manipulation, preparation of genomic DNAs, genetic analysis are performed, unless otherwise mentioned, in accordance with laboratory books, for example, "Experimental Protocols for Model Plants" (Shimamoto & Okada (1996), Cell Engineering SUPPLEMENT, Plant Cell Engineering Series 4, Shujunsha).

Example 1

Isolating *Arabidopsis* flowering regulating gene "MPC1"

[0045] In order to clone a flowering regulation gene from *Arabidopsis*, mutants that flower immediately after cotyledon expansion (super early flowering mutants) was isolated as follows. The M2 (seeds after self-fertilization of individuals obtained by sowing mutagenized seeds) of *Arabidopsis* (ecotype: *Landsberg*) mutagenized with chemical mutagen EMS were prepared and sown on agar medium (1/2 B5 medium (Gamborg et al. (1968), Exp. Cell Res., vol.50: p151-); 1% sucrose, 0.8% agar). The screening was performed by observing the morphology of seedlings germinated. About fifty thousand individuals divided into ten lots were screened to obtain one kind of super early flowering mutant. This mutant was named "mpc1." Since "mpc1" flowers before the plant sufficiently matures, its flower has no fertility and the strain cannot be maintained. Five thousand individuals from the lot in which the mutant had been obtained were cultivated to obtain seeds (M3) of each individual. Strains of heterozygotes that segregate the super early flowering mutant were obtained by sowing these seeds individually and observing their seedling. The "mpc1" mutation was found to be caused by single recessive gene from the facts that the mutation segregates the mutant individuals in the proportion of one fourth in the M3 generation and also segregates the mutant individuals in the proportion of one fourth in the F2 generation obtained by mating with the wild type.

[0046] The strains of heterozygotes were backcrossed onto wild type strain *Landsberg* two times and the next generation obtained was mated with wild type strain *Columbia*. A DNA was extracted from each individual of the F2 generation by the conventional methods and analyzed for the recombinant value between the mutant characteristics and RFLP (Restriction Fragment Length Polymorphism) marker (Liu et al. (1996), Plant J., vol.10(4): p733-), CAPS (Co-dominant cleaved Amplified Polymorphism Sequences) marker (Konienczmy et al. (1993), Plant J., vol.4: p403-), and microsatellite marker (Bell et al. (1994), Genomics, vol.19: p137-) to map the gene causing the mutation on a chromosome. The desired gene was mapped between well-known DNA markers on the chromosome 5, mi2 (Lister & Dean (1995), Weeds World, vol.2(1): p23-, <http://nasc.life.nott.ac.uk:8300/>) and Ds389-14 (Smith et al. (1996), Plant J., vol.10(4): p721-).

[0047] In order to isolate DNA fragments covering this chromosome region, C1C-YAC library (Creusot et al. (1995), Plant J., vol.8: p763-), P1-library (Liu et al. (1995), Plant J., vol.7: p351-) and TAC library (Liu et al. (1995), The Molecular Biology Society of Japan 18th Annual Meeting) were screened with the two marker mentioned above and DNA clones were obtained. DNA fragments were prepared from the clones obtained, novel DNA markers were generated, and detailed chromosome mapping of genes and screening of DNA clones were repeated on after another. As a result, the desired gene was found to locate between the markers 16EB53 and Z11-1, which can be obtained from the genomic DNA by PCR amplification. 16EB53 can be obtained by PCR with synthetic oligonucleotide primers "GGATCCGAAC CCGACTCGGT ACC" (SEQ ID NO: 4) and "GCTTATGGAT GTGGACTCTC TAAC" (SEQ ID NO: 5), and Z11-1 can be obtained by PCR with synthetic oligonucleotide primers "AGGTCCTACA ACTACAACAG TT" (SEQ ID NO: 6) and "GAGGAAGCTA GTATTCTCTT TG" (SEQ ID NO: 7).

[0048] The chromosome region between the markers 16EB53 and Z11-1 is indicated with DNA contigs of four kinds of TAC clones (11K22, 22K2, 19A20, and 20I12) shown in Figure 1. These TAC clones are about 70 to 100 kb long. When each of these clones was introduced into the mutant individual through *Agrobacterium tumefaciens*, the introduction of the three clones other than 20I12 reverted to wild-type (methods for gene introduction and cultivation of transformed plants are described in detail in Example 2). When cDNA library of *Arabidopsis* (Newman et al. (1994),

Plant Physiol., vol.106: p1241-) was screened using about 50 kb region common to these three clones as a probe, six kinds of gene cDNAs were obtained. The sites of these genes were mapped on DNA contigs. Moreover, the clones were completely or partially digested with restriction enzymes and subcloned to confirm whether each gene contributed to reversion. As a result of introducing these subclones, one gene having reversion ability was identified. This gene was confirmed to be the gene causing the super early flowering mutation "mpcl," that is, flowering regulating gene "MPC1". Analysis of this genomic region and the nucleotide sequence of the cDNA clones clarified that "MPC1" structural gene has 22 exons divided by 21 introns and that the length is 5580 bp. The protein encoded by the gene has a molecular weight of 69.5 kDa with 611 amino acid residues. It contains a C2H2 type zinc finger (Rosenfeld et al. (1993), J. Biomol. Struct. Dyn., vol.11: p557), which is characteristic of nucleic acid binding proteins, and an acidic amino acid cluster, in transcription activating domain of a transcription factor at amino acids 306 to 327 and 503 to 520, respectively, of SEQ ID NO: 1.

[0049] Homology search using DDBJ/EMBL/GenBank database detected sequences having partial homology, but each of them was a fragmentary sequence with unknown function. Specifically, they are a partial cDNA sequence of rice (EST C72616) and a genomic primary structure sequence of *Arabidopsis* (Z97342). This homologous sequence of *Arabidopsis* is located on the chromosome different from "MPC1" gene of the present invention and is greatly different from "MPC1" gene in that the region corresponding to that between the fifth and tenth exon of "MPC1" gene is missing. The sequence may be derived from the gene of the present invention by deletion of the above region, and thus be originally a gene related to flowering. These results indicates that genes homologous to that of the present invention with specific function have not been found so far and therefore the gene of the present invention is novel. In addition, the analysis of the nucleotide sequence of this gene of the "mpcl" mutant revealed that guanine base at 5039 of SEQ ID NO: 3 is replaced with adenine and that a termination codon occurs in the coding frame. An incomplete protein lacking amino acids from 541 and the following C-terminal region of "MPC1" protein by the base substitution is thought to be expressed in "mpcl" mutant. Since this protein lacks flowering regulating function partially or completely, it is thought that the plant cannot maintain vegetative growth and causes super early flowering.

Example 2

Inducing flowering by gene introduction

[0050] An antisense gene was constructed using a part of the *Arabidopsis* flowering regulating gene "MPC1" cDNA. The sequence between the BamHI site at nucleotide 1650 and the SphI site at 1984 of cDNA shown in SEQ ID NO: 2 was separated by restriction enzyme digestion. To transcribe the complementary sequence of the transcript of the resulting fragment, binary vector pBI121 (Jefferson et al. (1987), EMBO J., vol.6: p3901-) was cleaved at the XbaI site at the downstream of 35 S promoter, blunted, cleaved with BamHI, and ligated with the fragment obtained above that had been cleaved with SphI, blunted, and cleaved again with BamHI. This construct was used as an antisense gene.

[0051] The antisense gene was introduced into *Arabidopsis* by a gene transfer method using *Agrobacterium tumefaciens*. First, the antisense gene expression vector mentioned above was transferred into *Agrobacterium tumefaciens* by electroporation. The expression vector has the kanamycin resistance gene as the marker. The antisense gene expression vector DNA was mixed with *Agrobacterium tumefaciens* suspended in 10% glycerol and the mixture had electric pulse added in a 1 mm wide cuvette electrodes with a setting of 25 μ F, 600 Ω , and 1.8 kV. The cells were then cultivated on LB agar medium (1% bactotryptone, 0.5% yeast extract, 0.5% sodium chloride, 1.2% bactoagar) supplemented with 25 μ g/ml kanamycin and 50 μ g/ml rifampicin at 28°C for two days and colonies of kanamycin resistant *Agrobacterium tumefaciens* were screened. *Agrobacterium tumefaciens* having this antisense gene was cultivated in LB liquid medium (1% bactotryptone, 0.5% yeast extract, 0.5% sodium chloride) supplemented with 50 μ g/ml rifampicin and 25 μ g/ml kanamycin at 28°C for 18 hours to prepare culture of *Agrobacterium tumefaciens*.

[0052] Seeds of *Arabidopsis* sterilized with 1% sodium hypochlorite were sowed in MS agar medium (Murashige & Skoog (1962), Physiol. Plant, vol.15: p473-) supplemented with 1% sucrose and 0.4% Gellan Gum, and grown at 25°C for 14 days. A hypocotyl of grown *Arabidopsis* was cut out and put on CIM medium (B5 agar medium (Gamborg et al. (1968), Exp. Cell Res., vol.50: p151-) supplemented with 0.5 mg/l 2,4D, 0.05 mg/l kinetin, 2% glucose, and 0.4% Gellan Gum), and cultivated at 25°C for 6 days in the dark.

[0053] This hypocotyl was mixed with the above culture of *Agrobacterium tumefaciens* having the antisense gene, put on CIM medium again and cultivated at 25°C for two days to infect the hypocotyl with the bacteria. The hypocotyl was sterilized by washing in B5 liquid medium comprising 150 mg/l Claforan (cefotaxime sodium) and 2% glucose for five hours with shaking. The resulting hypocotyl was subcultured in SIM medium (B5 agar medium containing 2 mg/l gelatin, 0.2 mg/l IBA, 150 mg/l Claforan, 50 μ g/ml kanamycin, 2% glucose, and 0.4% Gellan Gum) every week to differentiate and screen the transformant. 35 S promoter, which promotes to express the antisense gene, is known as a constitutive expression promoter. Pistils were differentiated directly from the hypocotyl in fourth week of subcultivation.

Example 3Isolating rice flowering regulating gene "Os-MPC1"

5 [0054] Homology search of DDBJ/EMBL/GenBank database with the sequence of *Arabidopsis* flowering regulating gene "MPC1" as the probe detected a partial cDNA sequence of rice (EST C72616) that is partially homologous to MPC1. The whole cDNA sequence of this gene, whose function was unknown, was isolated as follows. First, cDNA library derived from rice immature seeds was screened using the partial cDNA sequence on the databases as the hybridization probe and one kind of cDNA clone was obtained. The cDNA of the clone was found to be 2248 bp long
 10 by nucleotide sequence determination and the protein encoded by this cDNA has a molecular weight of 68.6 kDa with 604 amino acid residues. The nucleotide sequence of the cDNA and the amino acid sequence of the protein encoded by the cDNA are shown in SEQ ID NO: 9 and 8, respectively. The sequence identity between the amino acid sequence of this protein and that of *Arabidopsis* "MPC1" protein is 61%, which is significant homology. Therefore, this gene was thought to be the gene corresponding to *Arabidopsis* "MPC1" in rice and was named "Os-MPC1." "Os-MPC1" protein
 15 was found to have a zinc finger motif and an acidic amino acid cluster as "MPC1" at amino acids 310 to 331 and 488 to 505, respectively, of SEQ ID NO: 8.

Example 420 Chromosome mapping of rice "Os-MPC1" gene

[0055] Chromosomal DNA fragments derived from a part of 3' region of "Os-MPC1" gene were amplified from rice strains "Asominori" and "IR24" by PCR. Synthetic oligonucleotide primers "GACGAGAAAC TTATTATGCA TATG" (SEQ ID NO: 10) and "GGTCTTGATA CTGCTCTACA GTTATG" (SEQ ID NO: 11) were used for amplification. About 1.3 kb
 25 amplified gene fragments thus obtained showed restriction fragment length polymorphism (RFLP); cleavage patterns were different between strains of, "Asominori" and "IR24" when digested with restriction enzyme SspI. The locus of "Os-MPC1" gene on a chromosome can be determined by correlating this RFLP with the RFLP map already known for rice. The locus of "Os-MPC1" was determined by calculating recombination value between "Os-MPC1" gene fragments and RFLP markers whose sites have been already determined using the chromosomal DNA of Recombinant
 30 Inbred (RI) strains generated from plants obtained by mating between rice "Asominori" and "IR24" (Tsunematsu et al. (1993), Rice Genetics Newsletter, vol.10: p89-). The result of the analysis revealed that "Os-MPC1" gene was located near the well-known C152 marker at the terminus of rice chromosome 9. Any flowering regulating genes have never been found at the terminus of the chromosome 9 of rice. From this fact, "Os-MPC1" gene is a novel and fundamental flowering regulating gene, which was difficult to be detected by conventional techniques.

Example 5Complementing *Arabidopsis* super early flowering mutation by rice "Os-MPC1" gene

40 [0056] Flowering regulating function of rice "Os-MPC1" gene isolated was tested. "Os-MPC1" cDNA was first cleaved at the NotI site at the 3'-terminal connection with the vector, blunted, and then cleaved at the NheI site in the 5' noncoding region to obtain only the cDNA sequence without the vector sequence. Separately, binary vector pBI121 (Jefferson et al. (1987), EMBO J., vol.6: p3901-) was cleaved at the SmaI site at the downstream of 35 S promoter and ligated with the blunted 3'-end of the above-mentioned cDNA fragment. The ligation product was then cleaved at the XbaI site at
 45 the upstream of the SmaI site in the vector, ligated at this site with the NheI site of the cDNA fragment to obtain the expression vector of "Os-MPC1" gene. The "Os-MPC1" gene was introduced into the mutant by introducing the above vector into *Agrobacterium tumefaciens* and infecting the slice of root of the *Arabidopsis* super early flowering mutant with the bacteria (methods for gene introduction and cultivation of plants into which the mutation was introduced are described in detail in Example 2). When a root of the super early flowering mutant without the "Os-MPC1" gene was
 50 cultivated to allow it to differentiate to an individual, only the direct floral differentiation due to the influence of the mutation was observed. In contrast, it was confirmed when "Os-MPC1" gene was introduced into the mutant that the mutation was complemented and stems and leaves were differentiated and grew.

[0057] These results indicate that not only "Os-MPC1" gene is functionally proved to be a flowering regulating gene of rice but also "flowering regulating gene" of the present invention functions similarly in wide-ranging species of plants.

55

Example 6Isolating flowering regulating genes from various kinds of plants

- 5 [0058] The amino acid sequences encoded by *Arabidopsis* "MPC1" gene and rice "Os-MPC1" gene were compared with each other and two regions were selected from similar amino acid sequences commonly found in both flowering regulating proteins. Specifically, one region is "Lys Arg Gln Phe Phe His Ser" (SEQ ID NO: 12) at amino acids 484 to 490 of SEQ ID NO: 1 and the other is "Trp Ala Cys Glu Ala Phe" (SEQ ID NO: 13) at amino acids 558 to 563 of SEQ ID NO: 1. Next, four kinds of synthetic oligonucleotide primers KR1 "AAGCGGCAAT TTTAYCAYTC" (SEQ ID NO: 14),
 10 KR2 "AAGCGGCAGT TCTAYCAYTC" (SEQ ID NO: 15), KR3 "AAGCGGCAGT TCTAYCAYAG" (SEQ ID NO: 16), and KR4 "AAGCGGCAAT TTTAYCAYAG" (SEQ ID NO: 17) were prepared based on the amino acid sequence of SEQ ID NO: 12, and two kinds of synthetic oligonucleotide primers WA1 "AATACCTCAC ANGCCCCA" (SEQ ID NO: 18) and WA2 "AATACTTCGC ANGCCCCA" (SEQ ID NO: 19) were prepared based on the amino acid sequence of SEQ ID NO: 13.
- 15 [0059] PCR was performed using eight kinds of combinations of the primers KR1, KR2, KR3, and KR4 with the primers WA1 and WA2 and chromosomal DNA of rice (ecotype: Nipponbare) and sugar beet (ecotype: Sugarman Gold) as templates. The nucleotide sequence of each amplified fragments was determined and compared to the known flowering regulating genes.
- 20 [0060] As a result, the 1216 bp fragment of sugar beet amplified by PCR using the primers KR1 and WA2 was proved to be a part of sugar beet flowering regulating gene. The nucleotide sequence of this sugar beet gene fragment is shown in SEQ ID NO: 20. In the gene fragment, nucleotide sequence encoding amino acids are divided by three introns and their locations are the same as that of *Arabidopsis* "MPC1" gene.
- [0061] The fragment amplified for rice by PCR using the primers KR2 and WA2 was proved to be a part of rice flowering regulating gene "Os-MPC1."
- 25 [0062] Using these amplified gene fragments, the full-length of the gene can be readily cloned by screening library clones, PCR technique, or other methods.
- [0063] It is possible to obtain flowering regulating genes from not only rice and sugar beet but also various species of plants using the method mentioned above.

Annex to the description

[0064]

5

SEQUENCE LISTING

10

<110> MITSUI CHEMISTRY CO., LTD.

<120> GENE FOR FLORAL REGURATION AND METHODS FOR CONTROLLING FLOWERING

15

<130> M4-101DP1

<140>

<141>

20

<150> JP 1998-180065

<151> 1998-06-26

25

<160> 22

<170> PatentIn Ver. 2.0

30

<210> 1

<211> 611

<212> PRT

35

<213> Arabidopsis thaliana

<220>

40

<221> ZN_FING

<222> (306)..(327)

45

<400> 1

Met Cys His Glu Asp Ser Arg Leu Arg Ile Ser Glu Glu Glu Glu Ile

1

5

10

15

50

Ala Ala Glu Glu Ser Leu Ala Ala Tyr Cys Lys Pro Val Glu Leu Tyr

20

25

30

55

Asn Ile Ile Gln Arg Arg Ala Ile Arg Asn Pro Leu Phe Leu Gln Arg

35

40

45

EP 0 967 278 A2

5	Cys Leu His Tyr Lys Ile Glu Ala Lys His Lys Arg Arg Ile Gln Met			
	50	55	60	
10	Thr Val-Phe Leu Ser Gly Ala Ile Asp Ala Gly Val Gln Thr Gln Lys			
	65	70	75	80
15	Leu Phe Pro Leu Tyr Ile Leu Leu Ala Arg Leu Val Ser Pro Lys Pro			
		85	90	95
20	Val Ala Glu Tyr Ser Ala Val Tyr Arg Phe Ser Arg Ala Cys Ile Leu			
		100	105	110
25	Thr Gly Gly Leu Gly Val Asp Gly Val Ser Gln Ala Gln Ala Asn Phe			
		115	120	125
30	Leu Leu Pro Asp Met Asn Arg Leu Ala Leu Glu Ala Lys Ser Gly Ser			
		130	135	140
35	Leu Ala Ile Leu Phe Ile Ser Phe Ala Gly Ala Gln Asn Ser Gln Phe			
		145	150	155
40	Gly Ile Asp Ser Gly Lys Ile His Ser Gly Asn Ile Gly Gly His Cys			
		165	170	175
45	Leu Trp Ser Lys Ile Pro Leu Gln Ser Leu Tyr Ala Ser Trp Gln Lys			
		180	185	190
50	Ser Pro Asn Met Asp Leu Gly Gln Arg Val Asp Thr Val Ser Leu Val			
		195	200	205
55	Glu Met Gln Pro Cys Phe Ile Lys Leu Lys Ser Met Ser Glu Glu Lys			
		210	215	220
	Cys Val Ser Ile Gln Val Pro Ser Asn Pro Leu Thr Ser Ser Ser Pro			
	225	230	235	240

EP 0 967 278 A2

5	Gln Gln Val Gln Val Thr Ile Ser Ala Glu Glu Val Gly Ser Thr Glu	245	250	255
10	Lys Ser Pro Tyr Ser Ser Phe Ser Tyr Asn Asp Ile Ser Ser Ser Ser	260	265	270
15	Leu Leu Gln Ile Ile Arg Leu Arg Thr Gly Asn Val Val Phe Asn Tyr	275	280	285
20	Arg Tyr Tyr Asn Asn Lys Leu Gln Lys Thr Glu Val Thr Glu Asp Phe	290	295	300
25	Ser Cys Pro Phe Cys Leu Val Lys Cys Ala Ser Phe Lys Gly Leu Arg	305	310	315
30	Tyr His Leu Pro Ser Thr His Asp Leu Leu Asn Phe Glu Phe Trp Val	325	330	335
35	Thr Glu Glu Phe Gln Ala Val Asn Val Ser Leu Lys Thr Glu Thr Met	340	345	350
40	Ile Ser Lys Val Asn Glu Asp Asp Val Asp Pro Lys Gln Gln Thr Phe	355	360	365
45	Phe Phe Ser Ser Lys Lys Phe Arg Arg Arg Arg Gln Lys Ser Gln Val	370	375	380
50	Arg Ser Ser Arg Gln Gly Pro His Leu Gly Leu Gly Cys Glu Val Leu	385	390	395
55	Asp Lys Thr Asp Asp Ala His Ser Val Arg Ser Glu Lys Ser Arg Ile	405	410	415
	Pro Pro Gly Lys His Tyr Glu Arg Ile Gly Gly Ala Glu Ser Gly Gln	420	425	430

EP 0 967 278 A2

5	Arg Val Pro Pro Gly Thr Ser Pro Ala Asp Val Gln Ser Cys Gly Asp	435	440	445
10	Pro Asp Tyr Val Gln Ser Ile Ala Gly Ser Thr Met Leu Gln Phe Ala	450	455	460
15	Lys Thr Arg Lys Ile Ser Ile Glu Arg Ser Asp Leu Arg Asn Arg Ser	465	470	475
20	Leu Leu Gln Lys Arg Gln Phe Phe His Ser His Arg Ala Gln Pro Met	485	490	495
25	Ala Leu Glu Gln Val Leu Ser Asp Arg Asp Ser Glu Asp Glu Val Asp	500	505	510
30	Asp Asp Val Ala Asp Phe Glu Asp Arg Arg Met Leu Asp Asp Phe Val	515	520	525
35	Asp Val Thr Lys Asp Glu Lys Gln Met Met His Met Trp Asn Ser Phe	530	535	540
40	Val Arg Lys Gln Arg Val Leu Ala Asp Gly His Ile Pro Trp Ala Cys	545	550	555
45	Glu Ala Phe Ser Arg Leu His Gly Pro Ile Met Val Arg Thr Pro His	565	570	575
50	Leu Ile Trp Cys Trp Arg Val Phe Met Val Lys Leu Trp Asn His Gly	580	585	590
55	Leu Leu Asp Ala Arg Thr Met Asn Asn Cys Asn Thr Phe Leu Glu Gln	595	600	605
	Leu Gln Ile	610		

```

5      <210> 2
      <211> 2280
      <212> DNA
      <213> Arabidopsis thaliana

10     <220>
      <221> CDS
      <222> (310)..(2142)

15     <220>
      <221> misc_feature
      <222> (1650)..(1655)
      <223> BamHI recognition site

20     <220>
      <221> misc_feature
      <222> (1984)..(1989)
      <223> SphI recognition site

25     <220>
      <221> misc_feature
      <222> (1984)..(1989)
      <223> SphI recognition site

30     <400> 2
      aagataattt ctcacaatta gggttttttt tttcttctga gttaaactgtt ccattctccat 60

      cctaattcttc accttctcct tgatttcgag atctctgtca attgttgaa tctgttcttt 120

40     atctaattag ctcaactccg agtctttgct ggattttgaa gctttttag ctgaagcaaa 180

      ttgttaatct gtgatggtgt atgcactgat tctgggtatg gtattgtact ctaggatctc 240

45     giagcgagaa tgccaggcat tcctcttggt agtcgtgaaa cctcttcttg ttcaagaagc 300

      acagagcag atg tgc cat gaa gac tcc cgt ctg cgt att tcg gaa gag gag 351
      Met Cys His Glu Asp Ser Arg Leu Arg Ile Ser Glu Glu Glu

50     1             5             10

55

```


EP 0 967 278 A2

gag att gct gct gaa gag agc ttg gct gcc tat tgc aag cct gtt gaa 399
 5 Glu Ile Ala Ala Glu Glu Ser Leu Ala Ala Tyr Cys Lys Pro Val Glu
 15 20 25 30

ctc tac-aat atc att caa cgc cgt gct att agg aat ccc ttg ttt ctt 447
 10 Leu Tyr Asn Ile Ile Gln Arg Arg Ala Ile Arg Asn Pro Leu Phe Leu
 35 40 45

cag cga tgt ttg cat tat aag att gag gca aaa cat aaa agg aga ata 495
 15 Gln Arg Cys Leu His Tyr Lys Ile Glu Ala Lys His Lys Arg Arg Ile
 50 55 60

caa atg act gta ttc ctc tcg ggc gct ata gat gct ggg gta caa act 543
 20 Gln Met Thr Val Phe Leu Ser Gly Ala Ile Asp Ala Gly Val Gln Thr
 65 70 75

caa aaa tta ttc cct ctg tat att ttg ttg gca aga ctc gtt tct cct 591
 30 Gln Lys Leu Phe Pro Leu Tyr Ile Leu Leu Ala Arg Leu Val Ser Pro
 80 85 90

aag cct gtc gct gag tat tct gca gta tat agg ttc agt cga gca tgt 639
 35 Lys Pro Val Ala Glu Tyr Ser Ala Val Tyr Arg Phe Ser Arg Ala Cys
 95 100 105 110

atc cta act ggt gga ttg ggg gtt gat gga gtt agt caa gcc caa gcc 687
 40 Ile Leu Thr Gly Gly Leu Gly Val Asp Gly Val Ser Gln Ala Gln Ala
 115 120 125

aac ttt ctt ctc cct gat atg aat aga ctc gca ttg gag gca aaa tca 735
 45 Asn Phe Leu Leu Pro Asp Met Asn Arg Leu Ala Leu Glu Ala Lys Ser
 130 135 140

gga tca ctc gct atc ttg ttt atc agc ttt gct ggt gcg caa aat tct 783
 50 Gly Ser Leu Ala Ile Leu Phe Ile Ser Phe Ala Gly Ala Gln Asn Ser
 145 150 155

EP 0 967 278 A2

5	caa ttt ggc att gat tca ggc aag att cat tca gga aat ata gga gga	831
	Gln Phe Gly Ile Asp Ser Gly Lys Ile His Ser Gly Asn Ile Gly Gly	
	160 165 170	
10	cat tgt tta tgg agc aaa ata cct ctg caa tca ctg tat gcg tcg tgg	879
	His Cys Leu Trp Ser Lys Ile Pro Leu Gln Ser Leu Tyr Ala Ser Trp	
	175 180 185 190	
15	cag aaa tca cca aac atg gac ttg gga cag aga gta gac aca gtc tct	927
	Gln Lys Ser Pro Asn Met Asp Leu Gly Gln Arg Val Asp Thr Val Ser	
	195 200 205	
20	ctt gtt gaa atg cag cct tgc ttc ata aag cta aag tcc atg agt gag	975
	Leu Val Glu Met Gln Pro Cys Phe Ile Lys Leu Lys Ser Met Ser Glu	
25	210 215 220	
30	gaa aag tgt gtc tcg att cag gtg ccc agc aat cca ctc acc tcg agc	1023
	Glu Lys Cys Val Ser Ile Gln Val Pro Ser Asn Pro Leu Thr Ser Ser	
	225 230 235	
35	tct ccg cag caa gtg caa gtc acc ata tct gca gaa gaa gtt ggg tca	1071
	Ser Pro Gln Gln Val Gln Val Thr Ile Ser Ala Glu Glu Val Gly Ser	
	240 245 250	
40	acg gaa aaa tct cct tat agt tca ttt tca tat aat gac atc tct tcc	1119
	Thr Glu Lys Ser Pro Tyr Ser Ser Phe Ser Tyr Asn Asp Ile Ser Ser	
	255 260 265 270	
45	tct tcc ttg ttg caa att atc agg ttg aga aca gga aat gta gtt ttc	1167
	Ser Ser Leu Leu Gln Ile Ile Arg Leu Arg Thr Gly Asn Val Val Phe	
50	275 280 285	
55	aac tac aga tac tat aac aac aaa ttg cag aag act gaa gta act gaa	1215
	Asn Tyr Arg Tyr Tyr Asn Asn Lys Leu Gln Lys Thr Glu Val Thr Glu	
	290 295 300	

EP 0 967 278 A2

5	gac ttt tct tgt cca ttc tgc tta gta aaa tgt gcc agt ttc aag ggc	1263
	Asp Phe Ser Cys Pro Phe Cys Leu Val Lys Cys Ala Ser Phe Lys Gly	
	305 310 315	
10	ctg aga tat cac ttg cca tca acc cac gat ctc ctc aat ttc gag ttt	1311
	Leu Arg Tyr His Leu Pro Ser Thr His Asp Leu Leu Asn Phe Glu Phe	
	320 325 330	
15	tgg gta act gaa gaa ttt cag gcg gta aat gtc tcc ctc aag act gag	1359
	Trp Val Thr Glu Glu Phe Gln Ala Val Asn Val Ser Leu Lys Thr Glu	
	335 340 345 350	
20	aca atg ata tcc aag gtt aat gag gat gac gtt gac cca aag cag caa	1407
	Thr Met Ile Ser Lys Val Asn Glu Asp Asp Val Asp Pro Lys Gln Gln	
	355 360 365	
25	act ttc ttt ttt tct tcc aaa aaa ttc aga cgg agg agg caa aag agt	1455
	Thr Phe Phe Phe Ser Ser Lys Lys Phe Arg Arg Arg Arg Gln Lys Ser	
30	370 375 380	
35	cag gta cgg agc tca agg caa ggg cct cat ctt gga tta ggt tgc gag	1503
	Gln Val Arg Ser Ser Arg Gln Gly Pro His Leu Gly Leu Gly Cys Glu	
	385 390 395	
40	gtg cta gat aag act gat gat gct cat tct gtt aga agt gag aag agc	1551
	Val Leu Asp Lys Thr Asp Asp Ala His Ser Val Arg Ser Glu Lys Ser	
	400 405 410	
45	cga ata cca cct gga aag cat tac gaa aga att ggg ggt gct gag tct	1599
	Arg Ile Pro Pro Gly Lys His Tyr Glu Arg Ile Gly Gly Ala Glu Ser	
	415 420 425 430	
50	ggt caa aga gtt cct cct ggc acg agt cct gca gac gtg caa tca tgt	1647
	Gly Gln Arg Val Pro Pro Gly Thr Ser Pro Ala Asp Val Gln Ser Cys	
55	435 440 445	

EP 0 967 278 A2

	ggg gat cca gat tat gtg cag tcg ata gct gga agt aca atg ttg cag	1695
5	Gly Asp Pro Asp Tyr Val Gln Ser Ile Ala Gly Ser Thr Met Leu Gln	
	450 455 460	
	ttt gca aaa acg agg aaa ata tct ata gaa cgg tcg gac ttg agg aac	1743
10	Phe Ala Lys Thr Arg Lys Ile Ser Ile Glu Arg Ser Asp Leu Arg Asn	
	465 470 475	
	cga agc ctc ctt cag aag aga cag ttc ttc cac tct cat cga gct cag	1791
15	Arg Ser Leu Leu Gln Lys Arg Gln Phe Phe His Ser His Arg Ala Gln	
	480 485 490	
	ccc atg gct cta gaa caa gta ctt tcg gac cgg gat agt gaa gat gaa	1839
20	Pro Met Ala Leu Glu Gln Val Leu Ser Asp Arg Asp Ser Glu Asp Glu	
	495 500 505 510	
25		
	gtt gat gat gat gtg gca gat ttt gaa gat aga agg atg ctc gat gat	1887
30	Val Asp Asp Asp Val Ala Asp Phe Glu Asp Arg Arg Met Leu Asp Asp	
	515 520 525	
	ttc gtt gat gtg act aaa gat gag aaa cag atg atg cac atg tgg aac	1935
35	Phe Val Asp Val Thr Lys Asp Glu Lys Gln Met Met His Met Trp Asn	
	530 535 540	
	tcg ttt gtg agg aag cag cga gta tta gca gat ggt cac att cca tgg	1983
40	Ser Phe Val Arg Lys Gln Arg Val Leu Ala Asp Gly His Ile Pro Trp	
	545 550 555	
	gca tgc gag gca ttc tca aga ttg cac gga ccc atc atg gtt cga aca	2031
45	Ala Cys Glu Ala Phe Ser Arg Leu His Gly Pro Ile Met Val Arg Thr	
	560 565 570	
50		
	ccg cac ttg att tgg tgc tgg aga gtg ttt atg gtg aaa ctg tgg aac	2079
55	Pro His Leu Ile Trp Cys Trp Arg Val Phe Met Val Lys Leu Trp Asn	
	575 580 585 590	

EP 0 967 278 A2

5 cac ggt ctt ctt gat gcc cga acc atg aac aac tgt aat acc ttt ctc 2127
 His Gly Leu Leu Asp Ala Arg Thr Met Asn Asn Cys Asn Thr Phe Leu
 595 600 605

10 gaa cag ctc caa att tgaaccacca agaatcatt aatttaagta gaaaaacasa 2182
 Glu Gln Leu Gln Ile
 610

15 gaaagacaag agaagaagag ttttgggttc tcatttaact acttttgggtg ttttaagaga 2242

20 aagaggagca tatttatgca tgaaaaaaaa aaaaaaaaa 2280

25 <210> 3
 <211> 5580
 <212> DNA
 <213> Arabidopsis thaliana

30 <220>
 <221> intron
 <222> (162)..(267)

35 <220>
 <221> intron
 <222> (394)..(565)

40 <220>
 <221> CDS
 <222> (588)..(713)

45 <220>
 <221> intron
 <222> (714)..(930)

50 <220>
 <221> CDS

55 <220>
 <221> CDS

	<222> (931)..(986)
5	<220>
	<221> intron
	<222> (987)..(1132)
10	<220>
	<221> CDS
15	<222> (1133)..(1247)
	<220>
	<221> intron
20	<222> (1248)..(1344)
	<220>
25	<221> CDS
	<222> (1345)..(1504)
	<220>
30	<221> intron
	<222> (1505)..(1596)
35	<220>
	<221> CDS
	<222> (1597)..(1647)
40	<220>
	<221> intron
45	<222> (1648)..(1854)
	<220>
	<221> CDS
50	<222> (1855)..(1994)
	<220>
55	<221> intron

<222> (1995)..(2197)

5

<220>

<221> CDS

<222> (2198)..(2260)

10

<220>

<221> intron

<222> (2261)..(2350)

15

<220>

<221> CDS

<222> (2351)..(2472)

20

<220>

<221> intron

<222> (2473)..(2714)

25

<220>

<221> CDS

<222> (2715)..(2779)

30

<220>

<221> intron

<222> (2780)..(2870)

35

<220>

<221> CDS

<222> (2871)..(2930)

40

<220>

<221> intron

<222> (2931)..(3038)

50

<220>

<221> CDS

55

	<222> (3039)..(3092)
5	<220>
	<221> intron
	<222> (3093)..(3174)
10	
	<220>
	<221> CDS
15	<222> (3175)..(3234)
	<220>
	<221> intron
20	<222> (3235)..(3654)
	<220>
25	<221> CDS
	<222> (3656)..(3701)
	<220>
30	<221> intron
	<222> (3702)..(3784)
35	<220>
	<221> CDS
	<222> (3785)..(3885)
40	
	<220>
	<221> intron
45	<222> (3886)..(4052)
	<220>
	<221> CDS
50	<222> (4053)..(4272)
	<220>
55	<221> intron

EP 0 967 278 A2

<222> (4273)..(4428)

5

<220>

<221> CDS

<222> (4429)..(4477)

10

<220>

<221> intron

<222> (4478)..(4552)

15

<220>

<221> CDS

20

<222> (4553)..(4636)

<220>

25

<221> intron

<222> (4637)..(4982)

<220>

30

<221> CDS

<222> (4983)..(5062)

35

<220>

<221> intron

<222> (5063)..(5265)

40

<220>

<221> CDS

45

<222> (5266)..(5355)

<220>

<221> intron

50

<222> (5356)..(5445)

<220>

55

<221> CDS

<222> (5446)..(5542)

5

<400> 3

aagataattt ctcacaatta gggttttttt ttcttctga gttactgtt ccctctccat 60

10

cctaattctt acccttctct tgatttcgag atctctgtca atttggtgaa tctgttcttt 120

atctaattag ctcaactccg agtctttgct ggattttgaa ggtcaccact gttcaagttt 180

15

acattttttt tctgtctaact cgttgatac cgtttctgc tgtgtggga tttattgggt 240

20

ttttcttctt tacgattttt gttgcagctt ttgtagctga agcaaatttg taatctgtga 300

tgggtgatgc actgattctg ggtatggtat tgtactctag gatctcgtag cgagaatgcc 360

25

aggcattcct ctgttagtc gtgaaacctc ttcgtaagtc tcatgaacaa cctaagctt 420

ctataatgtc tctgcagcat tgtgtaactt tatactgttt ctcttatgta taagctgagg 480

30

aatcctagta attcaaaactt atcaaatttt tattttgttg tggggttgct tacaattttg 540

35

gttgctgatg atggtgaaat cacagttggt caagaagcac agagcag atg tgc cat 596

Met Cys His

1

40

gaa gac tcc cgt ctg cgt att tcg gaa gag gag gag att gct gct gaa 644

Glu Asp Ser Arg Leu Arg Ile Ser Glu Glu Glu Glu Ile Ala Ala Glu

5

10

15

45

gag agc ttg gct gcc tat tgc aag cct gtt gaa ctc tac aat atc att 692

Glu Ser Leu Ala Ala Tyr Cys Lys Pro Val Glu Leu Tyr Asn Ile Ile

20

25

30

35

50

caa cgc cgt gct att agg aat gtatgtcttc ctccctacct ttttagaca gaat 747

Gln Arg Arg Ala Ile Arg Asn

55

40

EP 0 967 278 A2

atgtttagtt atgacttatg agctcagctg atatatcaca tgtattgggt tacttttgag 807

5 ttttgacaat gaaaatttac atgaaaatgt agtttgagtt gacttcattt ggtataagca 867

10 agtatgtgtt gtcttgctat gcagtcacac ctaatcattt ctctctctct gtctccctg 927

tag ccc ttg ttt ctt cag cga tgt ttg cat tat aag att gag gca aaa 975
Pro Leu Phe Leu Gln Arg Cys Leu His Tyr Lys Ile Glu Ala Lys

15 45 50 55

cat aaa agg ag gtaagctttt ttttttccct tcctttctct gttcagaatc tccatt 1032
His Lys Arg Arg

20 60

acttttgggt aactattaca ctatacctta gtaattcatt ccggacttga atgctttcta 1092

25 agttttcgga tagttatcaa tatatattac tgctttgcag a ata caa atg act gta 1148
Ile Gln Met Thr Val

30 65

ttc ctc tcg ggc gct ata gat gct ggg gta caa act caa aaa tta ttc 1196
Phe Leu Ser Gly Ala Ile Asp Ala Gly Val Gln Thr Gln Lys Leu Phe

35 70 75 80

cct ctg tat att ttg ttg gca aga ctc gtt tct cct aag cct gtc gct 1244
Pro Leu Tyr Ile Leu Leu Ala Arg Leu Val Ser Pro Lys Pro Val Ala

40 85 90 95

gag gtatgcattt gaacctcaga cagatttgca ttgatcttta ttatttgtaa cttacc 1303
Glu

45

tattctttgc taacattttt cttgaaattc tcaaattata g tat tct gca gta tat 1359
Tyr Ser Ala Val Tyr

50 100

55

EP 0 967 278 A2

5 agg ttc agt cga gca tgt atc cta act ggt gga ttg ggg gtt gat gga 1407
 Arg Phe Ser Arg Ala Cys Ile Leu Thr Gly Gly Leu Gly Val Asp Gly
 105 110 115 120

10 gtt agt caa gcc caa gcc aac ttt ctt ctc cct gat atg aat aga ctc 1455
 Val Ser Gln Ala Gln Ala Asn Phe Leu Leu Pro Asp Met Asn Arg Leu
 125 130 135

15 gca ttg gag gca aaa tca gga tca ctc gct atc ttg ttt atc agc ttt g 1504
 Ala Leu Glu Ala Lys Ser Gly Ser Leu Ala Ile Leu Phe Ile Ser Phe
 140 145 150

20 gtgattaaga ctgactgtgt acaaaattat ataaagacat ttatatatgt acagtattca 1564

25 gataaactga tcacataatt ttcttcttgt ag ct ggt gcg caa aat tct caa 1616
 Ala Gly Ala Gln Asn Ser Gln
 155

30 ttt ggc att gat tca ggc aag att cat tca g gtacttccat ttcttcattg a 1688
 Phe Gly Ile Asp Ser Gly Lys Ile His Ser
 160 165

35 tataacattc taatattgaa aagttatgta tctttgggca ttaccaattt tccatgtaat 1728

40 agtatggaaa atctcagtc tatttattaa caaaagaatt agggattctt tgactccaat 1788

 tataagagtt tctgaaagtc ttttttttca ttaactctta ccatcggaag cgtttttttc 1848

45 tgccag ga aat ata gga gga cat tgt tta tgg agc aaa ata cct ctg caa 1898
 Gly Asn Ile Gly Gly His Cys Leu Trp Ser Lys Ile Pro Leu Gln
 170 175 180

50 tca ctg tat gcg tgg tgg cag aaa tca cca aac atg gac ttg gga cag 1946
 Ser Leu Tyr Ala Ser Trp Gln Lys Ser Pro Asn Met Asp Leu Gly Gln
 55 185 190 195 200

EP 0 967 278 A2

aga gta gac aca gtc tct ctt gtt gaa atg cag cct tgc ttc ata aag g 1995
5 Arg Val Asp Thr Val Ser Leu Val Glu Met Gln Pro Cys Phe Ile Lys
205 210 215

taaacactat tgcccaagtc ttctcttgt tctatgactt tatgtccct gtattgaaat 2055
10 aaggactgtg tattgaactt ctttgttat ttgaaaaagt aaattggaag taattgctac 2115

tgtgaatttt atttttgcca ttagttttca gtcttgatta tttaaataaa aatattacgg 2175
15 tataacttgt ccattgctgc ag cta aag tcc atg agt gag gaa aag tgt gtc 2227
20 Leu Lys Ser Met Ser Glu Glu Lys Cys Val
220 225

tcg att cag gtg ccc agc aat cca ctc acc tcg gtaactttgc acactttgct 2280
25 Ser Ile Gln Val Pro Ser Asn Pro Leu Thr Ser
230 235

atacttccat acattattct gaaatatcat gtaatcatat tcttacaatt cttacacttc 2340
30 ttatttgaag agc tct ccg cag caa gtg caa gtc acc ata tct gca gaa 2389
35 Ser Ser Pro Gln Gln Val Gln Val Thr Ile Ser Ala Glu
240 245 250

gaa gtt ggg tca acg gaa aaa tct cct tat agt tca ttt tca tat aat 2437
40 Glu Val Gly Ser Thr Glu Lys Ser Pro Tyr Ser Ser Phe Ser Tyr Asn
255 260 265

gac atc tct tcc tct tcc ttg ttg caa att atc ag gtaactttca gttagt 2489
45 Asp Ile Ser Ser Ser Ser Leu Leu Gln Ile Ile Arg
270 275

ctgcaatttc ttctgcgtc tcagatttct tgccatcatc cattatgatt ttttgaatt 2549
50 gtataaaata tattggcgg tctgctatct cccttaatat atagttggca gttttcttga 2609
55

attgtgactg tcctcctctt ttatggggat tatacaagtc gttacgtaca actaaaaatg 2669
 5
 tccatctcgt taagttgact ctataccact acattcattg catag g ttg aga aca 2724
 Leu Arg Thr
 280
 10
 gga aat gta gtt ttc aac tac aga tac tat aac aac aaa ttg cag aag 2772
 Gly Asn Val Val Phe Asn Tyr Arg Tyr Tyr Asn Asn Lys Leu Gln Lys
 15
 285 290 295
 act gaa g gtaactagta ttattttaac ctgtttcata cccatgtgtg tctatatattc 2829
 20
 Thr Glu
 atccggtacc ctaacctgtt acgtatatgt ttgctatgtg tcttgacg ta act gaa 2885
 25
 Val Thr Glu
 300
 30
 gac ttt tct tgt cca ttc tgc tta gta aaa tgt gcc agt ttc aag gtgga 2935
 Asp Phe Ser Cys Pro Phe Cys Leu Val Lys Cys Ala Ser Phe Lys
 305 310 315
 35
 ctttcatttc cattctcatt catcctctta gtcaaagata cagctgtagt gactagtctt 2995
 40
 tgtagtgatg caatcttttc tttttctccc aatcatgttg tag ggc ctg aga tat 3050
 Gly Leu Arg Tyr
 320
 45
 cac ttg cca tca acc cac gat ctc ctc aat ttc gag ttt tgg gttgtagct 3101
 His Leu Pro Ser Thr His Asp Leu Leu Asn Phe Glu Phe Trp
 325 330 335
 50
 ttaaaattca gttaacctgt ttgatctttt ttttttattt tgtgggtgcc actaatctgc 3161
 55
 tttacttggg tag gta act gaa gaa ttt cag gcg gta aat gtc tcc ctc 3210

EP 0 967 278 A2

	Val Thr Glu Glu Phe Gln Ala Val Asn Val Ser Leu	
	340	345
5	aag act gag aca atg ata tcc aag gttagaacat cttgtttgtt cgatttatgt	3264
	Lys Thr Glu Thr Met Ile Ser Lys	
10	350 355	
	tcattagttt ctctgctgta tatttatag gctgtaacaa attcattttt catttaaact	3324
15	aatatcctcc atgggttggt gacttttgtg tggtaataa agggaactgg aatctttagt	3384
	tgctatttgt cacactatga tccttgctat tgccttaat agcgtgatga gaataaactc	3444
20	aaaatgacat cgtgttcig tttacttttt gtggccatga gaccgtcaaa gctcgactgt	3504
	agaataaagt cctggattat ataggagtgt caaatctaata tgaagtagtt ggttctacaa	3564
25	tatattctat gtctttgtag ttttccctat ttgatgatta ctcttagcac agttttctaa	3624
30	atgtaaatgt tcattaaaaa atctgctcag gtt aat gag gat gac gtt gac cca	3678
	Val Asn Glu Asp Asp Val Asp Pro	
	360	
35	aag cag caa act ttc ttt ttt tc gtaagtatac tggcctatat gtgcctttt	3731
	Lys Gln Gln Thr Phe Phe Phe Ser	
40	365 370	
	attatctttc cagcatctgt gtgagaccat aaaaattctt caatatgta cag t tcc	3788
45	Ser	
	aaa aaa ttc aga cgg agg agg caa aag agt cag gta cgg agc tca agg	3836
50	Lys Lys Phe Arg Arg Arg Arg Gln Lys Ser Gln Val Arg Ser Ser Arg	
	375 380 385	
55	caa ggg cct cat ctt gga tta ggt tgc gag gtg cta gat aag act gat g	3885

EP 0 967 278 A2

	Gln Gly Pro His Leu Gly Leu Gly Cys Glu Val Leu Asp Lys Thr Asp	
	390	395
		400
5		
	gatatgtgttt gactgaaatg acagttaatt ggattttag tagtggttc ttttgtgatg	3945
10	agagcctgtc ttagttgtat attttacgag tattttactt tgttatgtgc aattttgcat	4005
	gcaacaacgt tggatcattt ggcacagctt tttattctta ctttcag at gct cat	4060
15		Asp Ala His
		405
20	tct gtt aga agt gag aag agc cga ata cca cct gga aag cat tac gaa	4108
	Ser Val Arg Ser Glu Lys Ser Arg Ile Pro Pro Gly Lys His Tyr Glu	
	410	415
		420
25	aga att ggg ggt gct gag tct ggt caa aga gtt cct cct ggc acg agt	4156
	Arg Ile Gly Gly Ala Glu Ser Gly Gln Arg Val Pro Pro Gly Thr Ser	
	425	430
		435
30	cct gca gac gtg caa tca tgt ggg gat cca gat tat gtg cag tcg ata	4204
	Pro Ala Asp Val Gln Ser Cys Gly Asp Pro Asp Tyr Val Gln Ser Ile	
	440	445
		450
35		455
	gct gga agt acn atg ttg cag ttt gca aaa acg agg aaa ata tct ata	4252
	Ala Gly Ser Thr Met Leu Gln Phe Ala Lys Thr Arg Lys Ile Ser Ile	
40		460
		465
		470
45	gaa cgg tcg gac ttg agg aa gtatgttga cttccttttg tcgttctatc ctctt	4307
	Glu Arg Ser Asp Leu Arg Asn	
	475	
50	cttcaattta tatttaacta catatgggtc atgatgaaa aattgtgtcc tagttttata	4367
	acaagtagct tgtaaatccc aaatgatgtg agtgagtitt tcaaattttt tectctcca	4427
55	g c cga agc ctc ctt cag aag aga cag ttc ttc cac tct cat cga gct	4474

EP 0 967 278 A2

	Arg Ser Leu Leu Gln Lys Arg Gln Phe Phe His Ser His Arg Ala	
	480 485 490	
5		
	cag gtgatctttt ttcttttagct ctcttgccttt tgaagattgc aattgatttt gacttt	4533
	Gln	
10		
	gctatgtgta ctatcgag ccc atg gct cta gaa caa gta ctt tcg gac cgg	4585
	Pro Met Ala Leu Glu Gln Val Leu Ser Asp Arg	
15	495 500 505	
	gat agt gaa gat gaa gtt gat gat gat gtg gca gat ttt gaa gat aga	4633
20	Asp Ser Glu Asp Glu Val Asp Asp Asp Val Ala Asp Phe Glu Asp Arg	
	510 515 520	
25	agg gtatgttttt gaatttaata ttttcaccgc atcagtagtt gggtagaata aagctc	4692
	Arg	
30		
	agtagttggg tagatatacg tttcatgtga aagggaagg aatattgaag actgggcatg	4752
	ggcaaacggt aggagcaata ttgtaatggt tcagagatca atagaaaata tgtgagcaag	4812
35		
	cctcacggtt tgatatggaa cagtagaacc agatcattag tgcttatata acactcatta	4872
40	aaagacgaag tgtgtccggt tgtactcgat tctaacaatag ttgattctaa catagtttgt	4932
	ctgattctcc atatagtga taacgttatt tcctattact attctttcag atg ctc	4988
45	Met Leu	
	gat gat ttc gtt gat gtg act aaa gat gag aaa cag atg atg cac atg	5036
50	Asp Asp Phe Val Asp Val Thr Lys Asp Glu Lys Gln Met Met His Met	
	525 530 535 540	
55	tgg aac tcg ttt gtg agg aag cag cg gtatgtctta tctcttttca gtacatgt	5090

Trp Asn Ser Phe Val Arg Lys Gln Arg

545

5

cacgtggagt tttccagtat aaacatttag agtcgcgcat gtaaagggtg tggataattc 5150

10

ctgcctgggt tcttctggtt aaaaaaaaaa aactgaacaa ttagataaca tacgcatcca 5210

tgttctctga ctattataa gcattacctt gacagtgggt ttggaccctt tgcag a 5266

15

gta tta gca gat ggt cac att cca tgg gca tgc gag gca ttc tca aga 5314

Val Leu Ala Asp Gly His Ile Pro Trp Ala Cys Glu Ala Phe Ser Arg

550 555 560 565

20

ttg cac gga ccc atc atg gtt cga aca ccg cac ttg att tg gtaattcaac 5365

Leu His Gly Pro Ile Met Val Arg Thr Pro His Leu Ile Trp

25

570

575

tctcatttct tccattgttt tttccagtgt atcggagaag aaagcggttt tgttgataaa 5425

30

agtgagcttt ttttgtgtag g tgc tgg aga gtg ttt atg gtg aaa ctg tgg 5476

Cys Trp Arg Val Phe Met Val Lys Leu Trp

580

585

35

aac cac ggt ctt ctt gat gcc cga acc atg aac aac tgt aat acc ttt 5524

Asn His Gly Leu Leu Asp Ala Arg Thr Met Asn Asn Cys Asn Thr Phe

40

590

595

600

605

ctc gaa cag ctc caa att tgaaaaccca agaaatcatt aatttaagta gaaaaaca 5580

45

Leu Glu Gln Leu Gln Ile

610 611

50

<210> 4

<211> 23

<212> DNA

55

<213> Artificial Sequence

5 <220>
 <223> Description of Artificial Sequence:Artificially
 Synthesized Oligonucleotide Primer Sequence

 10 <400> 4
 ggatccgaac ccgactcggc acc 23

 15 <210> 5
 <211> 24
 <212> DNA
 20 <213> Artificial Sequence

 25 <220>
 <223> Description of Artificial Sequence:Artificially
 Synthesized Oligonucleotide Primer Sequence

 30 <400> 5
 gcttatggat gtggactctc taac 24

 35 <210> 6
 <211> 22
 <212> DNA
 40 <213> Artificial Sequence

 45 <220>
 <223> Description of Artificial Sequence:Artificially
 Synthesized Oligonucleotide Primer Sequence

 50 <400> 6
 aggtcctaca actacaacag tt 22

 55 <210> 7

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially
Synthesized Oligonucleotide Primer Sequence

<400> 7

gaggaagcta gtattctctt tg

22

<210> 8

<211> 604

<212> PRT

<213> Oryza sativa

<220>

<221> ZN_FING

<222> (310)..(335)

<400> 8

Met Cys Arg His Gln Pro Arg Ala Arg Leu Ser Pro Asp Glu Gln Leu

1 5 10 15

Ala Ala Glu Glu Ser Phe Ala Leu Tyr Cys Lys Pro Val Glu Leu Tyr

20 25 30

Asn Ile Ile Gln Arg Arg Ser Ile Lys Asn Pro Ala Phe Leu Gln Arg

35 40 45

Cys Leu Leu Tyr Lys Ile His Ala Arg Arg Lys Lys Arg Ser Leu Ile

50 55 60

Thr Ile Ser Leu Ser Gly Gly Thr Asn Lys Glu Leu Arg Ala Gln Asn

65 70 75 80

EP 0 967 278 A2

5	Ile Phe Pro Leu Tyr Val Leu Leu Ala Arg Pro Thr Asn Asn Val Ser	85	90	95	
10	Leu Glu Gly His Ser Pro Ile Tyr Arg Phe Ser Arg Ala Cys Leu Leu	100	105	110	
15	Thr Ser Phe His Glu Phe Gly Asn Lys Asp Tyr Thr Glu Ala Thr Phe	115	120	125	
20	Val Ile Pro Asp Val Lys Asn Leu Ala Thr Ser Arg Ala Cys Ser Leu	130	135	140	
25	Asn Ile Ile Leu Ile Ser Cys Gly Arg Ala Glu Gln Thr Phe Asp Asp	145	150	155	160
30	Asn Asn Cys Ser Gly Asn His Val Glu Gly Ser Thr Leu Gln Lys Leu	165	170	175	
35	Glu Gly Lys Cys Phe Trp Gly Lys Ile Pro Ile Asp Leu Leu Ala Ser	180	185	190	
40	Ser Leu Gly Asn Cys Val Ser Leu Ser Leu Gly His Thr Val Glu Met	195	200	205	
45	Ser Ser Thr Val Glu Met Thr Pro Ser Phe Leu Glu Pro Lys Phe Leu	210	215	220	
50	Glu Asp Asp Ser Cys Leu Thr Phe Cys Ser Gln Lys Val Asp Ala Thr	225	230	235	240
55	Gly Ser Phe Gln Leu Gln Val Ser Ile Ser Ala Gln Glu Ala Gly Ala	245	250	255	
	Lys Asp Met Ser Glu Ser Pro Tyr Ser Val Tyr Ser Tyr Asn Asp Val	260	265	270	

EP 0 967 278 A2

5	Pro Pro Ser Ser Leu Thr His Ile Ile Arg Leu Arg Ser Gly Asn Val	275	280	285
10	Leu Phe Asn Tyr Lys Tyr Tyr Asn Asn Thr Met Gln Lys Thr Glu Val	290	295	300
15	Thr Glu Asp Phe Ser Cys Pro Phe Cys Leu Val Pro Cys Gly Ser Phe	305	310	315
20	Lys Gly Leu Gly Cys His Leu Asn Ala Ser His Asp Leu Phe His Tyr	325	330	335
25	Glu Phe Trp Ile Ser Glu Glu Cys Gln Ala Val Asn Val Ser Leu Lys	340	345	350
30	Thr Asp Ser Trp Arg Thr Glu Leu Leu Ala Glu Gly Val Asp Pro Arg	355	360	365
35	His Gln Thr Phe Ser Tyr Arg Ser Arg Phe Lys Lys Arg Lys Arg Val	370	375	380
40	Glu Ile Ser Ser Asp Lys Ile Arg His Val His Pro His Ile Val Asp	385	390	395
45	Ser Gly Ser Pro Glu Asp Ala Gln Ala Gly Ser Glu Asp Asp Tyr Val	405	410	415
50	Gln Arg Glu Asn Gly Ser Ser Val Ala His Ala Ser Val Asp Pro Ala	420	425	430
55	Asn Ser Leu His Gly Ser Asn Leu Ser Ala Pro Thr Val Leu Gln Phe	435	440	445
	Gly Lys Thr Arg Lys Leu Ser Val Glu Arg Ala Asp Pro Arg Asn Arg	450	455	460

EP 0 967 278 A2

5 Gln Leu Leu Gln Lys Arg Gln Phe Phe His Ser His Arg Ala Gln Pro
465 470 475 480

10 Met Ala Trp Ser Lys Val Phe Ser Asp Arg Asp Ser Glu Asp Glu Val
485 490 495

15 Asp Asp Asp Ile Ala Asp Phe Glu Asp Arg Arg Met Leu Asp Asp Phe
500 505 510

20 Val Asp Val Thr Lys Asp Glu Lys Leu Ile Met His Met Trp Asn Ser
515 520 525

25 Phe Val Arg Lys Gln Arg Val Leu Ala Asp Gly His Ile Pro Trp Ala
530 535 540

30 Cys Glu Ala Phe Ser Gln Phe His Gly Gln Glu Leu Val Gln Asn Pro
545 550 555 560

35 Ala Leu Leu Trp Cys Trp Arg Phe Phe Met Val Lys Leu Trp Asn His
565 570 575

40 Ser Leu Leu Asp Ala Arg Ala Met Asn Ala Cys Asn Thr Ile Leu Glu
580 585 590

45 Gly Tyr Leu Asn Gly Ser Ser Asp Pro Lys Lys Asn
595 600

50 <210> 9
<211> 2248
<212> DNA
<213> Oryza sativa

55 <220>
<221> CDS

<222> (86)..(1897)

5

<220>

<221> misc_feature

<222> (36)..(41)

10

<223> NheI recognition site

<400> 9

15

cgccgatccc catccctccc gcgagcagga gcagggctag ccgtcgttcc tctgtgtgct 60

tccgccgcat ccctctgat accag atg tgc cgc cac cag cca agg gct cgg 112

Met Cys Arg His Gln Pro Arg Ala Arg

20

1

5

ctc tct ccc gat gag cag ctt gca gct gaa gaa agc ttc gca tta tac 160

25

Leu Ser Pro Asp Glu Gln Leu Ala Ala Glu Glu Ser Phe Ala Leu Tyr

10

15

20

25

tgc aag cag gtc gag ttg tat aat atc att cag cgc cga tcc att aaa 208

30

Cys Lys Pro Val Glu Leu Tyr Asn Ile Ile Gln Arg Arg Ser Ile Lys

30

35

40

aat cct gct ttt ctt caa aga tgc ctt ctt tac aag att cac gca aga 256

35

Asn Pro Ala Phe Leu Gln Arg Cys Leu Leu Tyr Lys Ile His Ala Arg

45

50

55

cgg aag aag agg agc ctg ata acc ata tca ctt tct gga ggc aca aat 304

40

Arg Lys Lys Arg Ser Leu Ile Thr Ile Ser Leu Ser Gly Gly Thr Asn

60

65

70

aaa gaa ctg cgg gca caa aat atc ttt cct ctt tat gtt ctg tta gct 352

45

Lys Glu Leu Arg Ala Gln Asn Ile Phe Pro Leu Tyr Val Leu Leu Ala

50

75

80

85

aga cct act aat aat gtt tca ctt gaa ggg cat tct ccg ata tat cga 400

55

Arg Pro Thr Asn Asn Val Ser Leu Glu Gly His Ser Pro Ile Tyr Arg

EP 0 967 278 A2

	90	95	100	105	
5	ttc agt cgt gct tgt ttg ttg act tct ttt cat gaa ttt gga aat aaa				448
	Phe Ser Arg Ala Cys Leu Leu Thr Ser Phe His Glu Phe Gly Asn Lys				
		110	115	120	
10	gac tac act gaa gca aca ttc gtc att cct gat gtg aag aac tta gca				496
	Asp Tyr Thr Glu Ala Thr Phe Val Ile Pro Asp Val Lys Asn Leu Ala				
15		125	130	135	
	acc tcc cga gct tgc agc ctt aat att atc ctt atc agc tgt gga cga				544
	Thr Ser Arg Ala Cys Ser Leu Asn Ile Ile Leu Ile Ser Cys Gly Arg				
20		140	145	150	
	gct gag caa act ttt gat gac aat aac tgt tct ggg aac cat gtg gaa				592
25	Ala Glu Gln Thr Phe Asp Asp Asn Asn Cys Ser Gly Asn His Val Glu				
		155	160	165	
	ggc tct act ctc caa aag ctt gaa ggg aag tgt ttc tgg ggt aaa ata				640
30	Gly Ser Thr Leu Gln Lys Leu Glu Gly Lys Cys Phe Trp Gly Lys Ile				
		170	175	180	185
	cca atc gat ctt ctt gct tca tct ttg gga aat tgt gtg agc tta agt				688
35	Pro Ile Asp Leu Leu Ala Ser Ser Leu Gly Asn Cys Val Ser Leu Ser				
		190	195	200	
40	ttg gga cat acc gtg gaa atg tct tcc acg gtt gag atg acc cca agc				736
	Leu Gly His Thr Val Glu Met Ser Ser Thr Val Glu Met Thr Pro Ser				
		205	210	215	
45	ttc tta gag cca aaa ttt ctg gag gat gac agt tgc ttg aca ttt tgc				784
	Phe Leu Glu Pro Lys Phe Leu Glu Asp Asp Ser Cys Leu Thr Phe Cys				
50		220	225	230	
	tct cag aag gtt gat gct act ggt tca ttt caa ctg caa gtt agc ata				832
55	Ser Gln Lys Val Asp Ala Thr Gly Ser Phe Gln Leu Gln Val Ser Ile				

EP 0 967 278 A2

	235	240	245	
5	tct gct caa gag gct ggt gca aaa gac atg tcc gag tct cct tat agt			880
	Ser Ala Gln Glu Ala Gly Ala Lys Asp Met Ser Glu Ser Pro Tyr Ser			
	250	255	260	265
10	gtt tat tca tat aat gat gtg cca cct tcg tca ttg aca cat att ata			928
	Val Tyr Ser Tyr Asn Asp Val Pro Pro Ser Ser Leu Thr His Ile Ile			
15		270	275	280
	agg ttg aga tct ggc aat gtg ctt ttt aac tac aaa tac tac aat aat			976
	Arg Leu Arg Ser Gly Asn Val Leu Phe Asn Tyr Lys Tyr Tyr Asn Asn			
20		285	290	295
	act atg caa aaa acc gaa gtc act gaa gat ttt tct tgc cca ttt tgc			1024
25	Thr Met Gln Lys Thr Glu Val Thr Glu Asp Phe Ser Cys Pro Phe Cys			
	300	305	310	
30	ttg gta cca tgt ggc agc ttt aag ggt cta gga tgt cac cta aac gca			1072
	Leu Val Pro Cys Gly Ser Phe Lys Gly Leu Gly Cys His Leu Asn Ala			
	315	320	325	
35	tcg cat gac ctt ttc cat tat gag ttt tgg ata tct gaa gag tgc cag			1120
	Ser His Asp Leu Phe His Tyr Glu Phe Trp Ile Ser Glu Glu Cys Gln			
	330	335	340	345
40	gct gtt aat gtt agt ctg aag act gat tct tgg aga aca gag ctt ttg			1168
	Ala Val Asn Val Ser Leu Lys Thr Asp Ser Trp Arg Thr Glu Leu Leu			
		350	355	360
45	gct gag gga gtt gat cca aga cat caa aca ttt tcg tac cgc tca aga			1216
	Ala Glu Gly Val Asp Pro Arg His Gln Thr Phe Ser Tyr Arg Ser Arg			
50		365	370	375
	ttt aag aag cgt aaa agg gtg gaa atc tca agt gat aaa att agg cat			1264
55	Phe Lys Lys Arg Lys Arg Val Glu Ile Ser Ser Asp Lys Ile Arg His			

EP 0 967 278 A2

	380	385	390	
5	gta cat cca cat att gtg gat tca gga tca cct gaa gat gcc cag gca	1312		
	Val His Pro His Ile Val Asp Ser Gly Ser Pro Glu Asp Ala Gln Ala			
	395	400	405	
10	gga tct gaa gac gat tac gtg cag agg gaa aat ggt agt tct gta gca	1360		
	Gly Ser Glu Asp Asp Tyr Val Gln Arg Glu Asn Gly Ser Ser Val Ala			
15	410	415	420	425
	cac gct tct gtt gat cct gct aat tca tta cac ggt agc aat ctt tca	1408		
	His Ala Ser Val Asp Pro Ala Asn Ser Leu His Gly Ser Asn Leu Ser			
20		430	435	440
	gca cca aca gtg tta cag ttt ggg aag aca aga aag ctg tct gtt gaa	1456		
25	Ala Pro Thr Val Leu Gln Phe Gly Lys Thr Arg Lys Leu Ser Val Glu			
	445	450	455	
	cga gct gat ccc aga aat cgg cag ctc cta caa aaa cgc cag ttc ttt	1504		
30	Arg Ala Asp Pro Arg Asn Arg Gln Leu Leu Gln Lys Arg Gln Phe Phe			
	460	465	470	
35	cat tct cac agg gct caa cca atg gca tgg agc aaa gtt ttc tca gat	1552		
	His Ser His Arg Ala Gln Pro Met Ala Trp Ser Lys Val Phe Ser Asp			
	475	480	485	
40	cgt gat agt gaa gat gaa gtt gat gat gac att gct gat ttt gaa gat	1600		
	Arg Asp Ser Glu Asp Glu Val Asp Asp Asp Ile Ala Asp Phe Glu Asp			
45	490	495	500	505
	aga aga atg ctt gat gat ttt gtt gat gtt aca aaa gac gag aaa ctt	1648		
	Arg Arg Met Leu Asp Asp Phe Val Asp Val Thr Lys Asp Glu Lys Leu			
50		510	515	520
	att atg cat atg tgg aat tca ttt gtt cgg aaa caa agg gta cta gcg	1696		
55	Ile Met His Met Trp Asn Ser Phe Val Arg Lys Gln Arg Val Leu Ala			

EP 0 967 278 A2

	525	530	535	
5	gat ggc cat att ccc tgg gca tgc gaa gca ttc tcg cag ttt cat gga			1744
	Asp Gly His Ile Pro Trp Ala Cys Glu Ala Phe Ser Gln Phe His Gly			
	-540	545	550	
10	caa gaa ctt gta caa aat cca gct cta cta tgg tgt tgg agg ttt ttt			1792
	Gln Glu Leu Val Gln Asn Pro Ala Leu Leu Trp Cys Trp Arg Phe Phe			
15	555	560	565	
20	atg gtc aaa ctc tgg aac cac agt cta ctg gat gcg cga gcc atg aat			1840
	Met Val Lys Leu Trp Asn His Ser Leu Leu Asp Ala Arg Ala Met Asn			
	570	575	580	585
25	gcc tgc aac aca att ctt gaa ggc tac ctg aac gga agc tcg gat cca			1888
	Ala Cys Asn Thr Ile Leu Glu Gly Tyr Leu Asn Gly Ser Ser Asp Pro			
	590	595	600	
30	aag aaa aat tgacgcatac aaatcattgg ccaacctgta gagtaaaatg			1937
	Lys Lys Asn			
35	cacttgctact ggctctggcc attccaatag ttgtttttgt ttttgaaaa aaagatgtct			1997
	gaagaattga aagctaacat gtgttttgga gggaagaaaa ttgaaggctg gggcggtcat			2057
40	tgtttcattt agaactcttc tcgattctat ttattgtaat tgatgttact cataactgta			2117
	gagcagtatc aagaccaaac tgtaatgata tggtagcaa tatttacata aaagtttatt			2177
45	ttgtttgttg tttagcaccg tgggcagaca atttaattcc tatgcaggcc ctttttcac			2237
50	gtcaaaaaaa a			2248
55	<210> 10			
	<211> 24			

5 <212> DNA
 <213> Artificial Sequence

 10 <220>
 <223> Description of Artificial Sequence:Artificially
 Synthesized Oligonucleotide Primer Sequence

 15 <400> 10
 gacgagaaac ttattatgca tatg 24

 20 <210> 11
 <211> 26
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> Description of Artificial Sequence:Artificially
 Synthesized Oligonucleotide Primer Sequence
 30
 <400> 11
 ggtcttgata ctgctctaca gttatg 26
 35
 <210> 12
 <211> 7
 <212> PRT
 <213> Arabidopsis thaliana
 45
 <400> 12
 Lys Arg Gln Phe Phe His Ser
 1 5
 50
 <210> 13
 <211> 6
 55

<212> PRT

<213> Arabidopsis thaliana

<400> 13

Trp Ala Cys Glu Ala Phe

1 5

<210> 14

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially
Synthesized Oligonucleotide Primer Sequence

<400> 14

aagcggaat ttatcaytc

20

<210> 15

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially
Synthesized Oligonucleotide Primer Sequence

<400> 15

aagcggaat ttatcaytc

20

<210> 16

<211> 20

5 <212> DNA
 <213> Artificial Sequence

 <220>
 10 <223> Description of Artificial Sequence:Artificially
 Synthesized Oligonucleotide Primer Sequence

 <400> 16
 15 aagcggcagt tctaycayag 20

 <210> 17
 20 <211> 20
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> Description of Artificial Sequence:Artificially
 30 Synthesized Oligonucleotide Primer Sequence

 <400> 17
 35 aagcggcaat ttaycayag 20

 <210> 18
 40 <211> 17
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> Description of Artificial Sequence:Artificially
 Synthesized Oligonucleotide Primer Sequence
 50
 <400> 18
 55 aatacctcac angccca 17

5 <210> 19
 <211> 17
 <212> DNA
 <213> Artificial Sequence
 10 <220>
 <223> Description of Artificial Sequence:Artificially
 Synthesized Oligonucleotide Primer Sequence
 15
 <400> 19
 aatacttcgc angccca
 20
 <210> 20
 25 <211> 1216
 <212> DNA
 <213> Beta vulgaris
 30 <220>
 <221> misc_feature
 <222> (1)..(20)
 35 <223> Oligonucleotide Primer "KR1" Sequence
 <220>
 40 <221> CDS
 <222> (21)..(33)
 45 <220>
 <221> intron
 <222> (34)..(694)
 50 <220>
 <221> CDS
 55 <222> (695)..(778)

17

EP 0 967 278 A2

<220>
 <221> intron
 5 <222> (779)..(951)

 <220>
 10 <221> CDS
 <222> (952)..(1031)

 <220>
 15 <221> intron
 <222> (1032)..(1174)

 <220>
 20 <221> CDS
 <222> (1175)..(1199)
 25
 <220>
 <221> misc_feature
 30 <222> (1200)..(1216)
 <223> Oligonucleotide Primer "WAZ" Sequence

 <400> 20
 35 aagcggcaat tttatcattc t cac aga gct cag gtaatcaact gcagaagtca tat 56
 His Arg Ala Gln
 1
 40
 cgtgttatgc tgatgtctga actcctataa tataacagtt gttgactctt tgtttcctat 116
 45 agtagttgtc ttgatggttg atcaaatttt gacaacattt cagcattctt aaacatcttt 176
 tcattatttt ttatttaca agagtagtaa ttcaagcacc ataagaaaca ctgatcaata 236
 50 gtttcttgca agttcttgaa cacttaataa gcagaggggt acttttaaatt attcagcatt 296
 55 tgtttgataa tctcaggtgt ttggacttg ctatatgtac ctgatgacac cgcttttagtt 356

EP 0 967 278 A2

tcaactagga tatggcgcta aatgggggaa aattgataaa gtcgagtagc aaaaatgatt 416

5 aggatttttaa cgtgggtgtt ctccttttct ctctcaagtt cattgtggg tgccatctat 476

agaaatgtct cgggttgtag ttttctatg gaaatgcagg cgtcgtttca gagtttgctc 536

10 tctgcttctc tcaatagtca attcagataa gccactttca ctgcaacctt gactgctact 596

cttggacttc aaattctagt cctctttgtc ttgtatcat tcttcaattt ttccaattga 656

15 tgatgctgat ttgaaaaac tctcttttgc acccgaag cca atg gct ctg gat caa 712

Pro Met Ala Leu Asp Gln

20 5 10

gta ttg tca gac agg gat agt gag gat gaa gtg gat gat gat att gct 760

25 Val Leu Ser Asp Arg Asp Ser Glu Asp Glu Val Asp Asp Asp Ile Ala

15 20 25

gct ctt gaa gat aga agg gtacgtttgg ttattttcca aattttttga gttgcttg 816

Ala Leu Glu Asp Arg Arg

30

35 cgtgattaac aatttttgat ctagtaatgg ttcttgcttc tagccaagtc ttgaatttc 876

taatglaata gttatctttt tcttgagtc attttgctaa ctaaaccgtg tatggtacct 936

40 tgccttgtgc tgcag atg ctt gat gat ttt gtg gat gta agc aaa gac gaa 987

Met Leu Asp Asp Phe Val Asp Val Ser Lys Asp Glu

45 35 40

aaa cac cta atg cat cta tgg aac tca ttt gta aaa aag caa ag gtagac 1037

50 Lys His Leu Met His Leu Trp Asn Ser Phe Val Lys Lys Gln Arg

45 50 55

55 ttgttatgc aattgtcccg ttgtttaat ttctttctcc attgtgaatg cttgcgtagt 1097

gtgctccgga agtatatttg atggcgctta cctgtggttg ttggctttg tgtaatgttt 1157

5 ccatttttgt gcaccag g gtt ttg gct gat ggt cat gtt ccc tgggcatgcg a 1211

Val Leu Ala Asp Gly His Val Pro

60 65 67

10 agtatt 1216

15 Claims

1. A DNA encoding a protein having flowering regulating activity, wherein said DNA selected from the group consisting of:
 - 20 i) a DNA encoding a protein comprising the amino acid sequence of SEQ ID NO: 1;
 - ii) a DNA encoding a protein comprising the amino acid sequence substantially identical to that of SEQ ID NO: 1;
 - iii) a DNA hybridizing a DNA encoding the protein comprising the amino acid sequence of SEQ ID NO:1;
 - iv) a DNA encoding a protein comprising amino acid sequences showing 50% or more and 60% or more homology with amino acids 278 to 348 and 465 to 607, respectively, of the amino acid sequence of SEQ ID NO: 1;
 - 25 v) a DNA encoding a protein comprising the amino acid sequence of SEQ ID NO: 8.
 - vi) a DNA encoding a protein comprising the amino acid sequence substantially identical to that of SEQ ID NO: 8.
 - vii) a DNA hybridizing a DNA encoding the protein comprising the amino acid sequence of SEQ ID NO: 8; and
 - 30 viii) a DNA encoding a protein comprising amino acid sequences showing 50% or more and 60% or more homology with amino acids 282 to 352 and 450 to 592, respectively, of the amino acid sequence of SEQ ID NO: 8.
2. The DNA of claim 1, wherein said DNA of i) comprises the coding region of the nucleotide sequence of SEQ ID NO: 2.
3. The DNA of claim 1, wherein said DNA of v) comprises the coding region of the nucleotide sequence of SEQ ID NO: 9.
4. The DNA of any one of claims 1 to 3, encoding a protein having a zinc finger structure.
5. A protein having flowering regulating activity, encoded by the DNA of any one of claims 1 to 4.
6. The protein of claim 5, comprising the amino acid sequence of SEQ ID NO: 1 or 8.
7. A recombinant double-stranded DNA molecule comprising an expression cassette comprising the DNA of any one of claims 1 to 4.
8. A recombinant double-stranded DNA molecule comprising an expression cassette comprising the following constituent elements of i) to iii),
 - i) a promoter that can transcribe in plant cells,
 - ii) the DNA of any one of claims 1 to 4 or a part of it fused to said promoter in sense or antisense direction, and selectively, and
 - 55 iii) a signal involved in transcription termination of RNA molecules and polyadenylation, wherein the signal functions in plants.
9. A transformant into which the recombinant double-stranded DNA molecule of claim 7 is introduced.

10. A transgenic plant cell into which the recombinant double-stranded DNA molecule of claim 8 is introduced.
11. A method for producing a protein of claim 5 or 6, wherein the method comprises (a) cultivating a transformant of claim 9 and (b) recovering a recombinant protein from said transformant or the culture supernatant of it.
- 5 12. A transgenic plant comprising transgenic plant cells of claim 10.
13. A method for producing a transgenic plant of claim 12, wherein said method comprises
- 10 (a) introducing the recombinant double-stranded DNA molecule of claim 8 into plant cells and
(b) regenerating said plant cells.
14. A DNA encoding an antisense RNA complementary to a transcription product of a DNA of any one of claims 1 to 4.
- 15 15. A method for regulating the flowering time of a plant, wherein said method comprises introducing the whole or a part of a DNA of any one of claims 1 to 4 or the whole or a part of a DNA of claim 14 into a plant and expressing it, thereby changing the activity of a flowering regulating protein.
- 20 16. An antibody that binds to a protein of claim 5 or 6.

Fig. 1

